


FORM PTO-1390 (REV.11-2000)	U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER 249-255
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		U.S. APPLICATION NO. (If known, see 37 C.F.R. 1.5) 10/089500 Unknown
INTERNATIONAL APPLICATION NO. PCT/JP00/06774	INTERNATIONAL FILING DATE 29 September 2000	PRIORITY DATE CLAIMED 30 September 1999
TITLE OF INVENTION COMPLEMENTARY DETERMINING REGION-GRAFTED ANTIBODY AGAINST GANGLIOSIDE GD3 AND DERIVATIVE OF ANTIBODY AGAINST GANGLIOSIDE GD3		
APPLICANT(S) FOR DO/EO/US HANAI et al		
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:		
1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input checked="" type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below. 4. <input checked="" type="checkbox"/> The U.S. has been elected by the expiration of 19 months from the priority date (Article 31). 5. A copy of the International Application as filed (35 U.S.C. 371(c)(2)). a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau). b. <input checked="" type="checkbox"/> has been communicated by the International Bureau. c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input checked="" type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). a. <input checked="" type="checkbox"/> is attached hereto. b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4). 7. <input type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)). a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> have been communicated by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> A English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). Items 11 To 20 below concern document(s) or information included: 11. <input type="checkbox"/> An Information Disclosure Statement under 37 C.F.R. 1.97 and 1.98. 12. <input checked="" type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 C.F.R. 3.28 and 3.31 is included. 13. <input type="checkbox"/> A FIRST preliminary amendment. 14. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 15. <input type="checkbox"/> A substitute specification. 16. <input type="checkbox"/> A change of power of attorney and/or address letter. 17. <input checked="" type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821-1.825. 18. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4). 19. <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4). 20. <input checked="" type="checkbox"/> Other items or information. PTO -1449 and copy of International Search Report and a paper copy of Sequence Listing; Budapest Treaty Deposit receipts relating to FERM BP-6918, FERM BP-6791 & FERM BP6790		

U.S. APPLICATION NO. (If known, see 37 C.F.R. 1.53) Unknown 107089500				INTERNATIONAL APPLICATION NO. PCT/JP00/06774				ATTORNEY'S DOCKET NUMBER 249-255					
21. <input checked="" type="checkbox"/> The following fees are submitted:								CALCULATIONS PTO USE ONLY					
BASIC NATIONAL FEE (37 C.F.R. 1.492(a)(1)-(5)): -- Neither international preliminary examination fee (37 C.F.R. 1.482) nor international search fee (37 C.F.R. 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO\$1040.00 -- International preliminary examination fee (37 C.F.R. 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO\$890.00 -- International preliminary examination fee (37 C.F.R. 1.482) not paid to USPTO but international search fee (37 C.F.R. 1.445(a)(2)) paid to USPTO\$740.00 -- International preliminary examination fee (37 C.F.R. 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4)\$710.00 -- International preliminary examination fee (37 C.F.R. 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4)\$100.00 <div style="text-align: right;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div>								\$		890.00			
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 C.F.R. 1.492(e)).								\$		0.00			
CLAIMS		NUMBER FILED		NUMBER EXTRA		RATE							
Total Claims		66		-20 =		46		X		\$18.00		\$ 828.00	
Independent Claims		2		-3 =		0		X		\$84.00		0.00	
MULTIPLE DEPENDENT CLAIMS(S) (if applicable)										\$280.00		\$ 0.00	
TOTAL OF ABOVE CALCULATIONS =												\$ 1718.00	
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.												0.00	
SUBTOTAL =												\$ 1718.00	
Processing fee of \$130.00, for furnishing the English Translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 C.F.R. 1.492(f)).												0.00	
TOTAL NATIONAL FEE =												\$ 1718.00	
Fee for recording the enclosed assignment (37 C.F.R. 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 C.F.R. 3.28, 3.31). \$40.00 per property												40.00	
Fee for Petition to Revive Unintentionally Abandoned Application (\$1280.00 - Small Entity = \$640.00)												0.00	
TOTAL FEES ENCLOSED =												\$ 1758.00	
								Amount to be:					
								refunded				\$	
								Charged				\$	
a. <input checked="" type="checkbox"/> A check in the amount of \$1758.00 to cover the above fees is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. 14-1140 in the amount of \$_____ to cover the above fees. A duplicate copy of this form is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>14-1140</u> . A duplicate copy of this form is enclosed. d. <input checked="" type="checkbox"/> The entire content of the foreign application(s), referred to in this application is/are hereby incorporated by reference in this application.													
NOTE: Where an appropriate time limit under 37 C.F.R. 1.494 or 1.495 has not been met, a petition to revive (37 C.F.R. 1.137(a) or (b)) must be filed and granted to restore the application to pending status.													
SEND ALL CORRESPONDENCE TO: NIXON & VANDERHYE P.C. 1100 North Glebe Road, 8 th Floor Arlington, Virginia 22201-4714 Telephone: (703) 816-4000													
										 SIGNATURE			
										B. J. Sadoff NAME			
										36,663 REGISTRATION NUMBER			
										March 29, 2002 Date			

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of

HANAI et al

Atty. Ref.: **249-255**

Serial No. **Unknown**

Group:

National Phase of: **PCT/JP00/06774**

International Filing Date: **29 September 2000**

Filed: **March 29, 2002**

Examiner:

For: **COMPLEMENTARY DETERMINING REGION-GRAFTED
ANTIBODY AGAINST GANGLIOSIDE GD3 AND
DERIVATIVE OF ANTIBODY AGAINST GANGLIOSIDE
GD3**

* * * * *

March 29, 2002

Assistant Commissioner for Patents
Washington, DC 20231

Sir:

PRELIMINARY AMENDMENT

Prior to calculation of the filing fee and in order to place the above identified application in better condition for examination, please amend as follows:

IN THE SPECIFICATION

Page 1, after the title insert the following:

-- This application is the US national phase of international application PCT/JP00/06774 filed September 29, 2000 which designated the U.S. --.

Insert the attached Sequence Listing in place of the originally-filed Sequence Listing.

IN THE CLAIMS

Please substitute the following amended claims for corresponding claims previously presented. A copy of the amended claims showing current revisions is attached.

HANAI et al
Serial No. **Unknown**

3. (Amended) The derivative of an antibody according to claim 1, wherein the monoclonal antibody comprises CDR1, CDR2 and CDR3 of H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

4. (Amended) The derivative of an antibody according to claim 1, wherein the monoclonal antibody comprises CDR1, CDR2 and CDR3 of L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

5. (Amended) The derivative of an antibody according to claim 1, wherein the monoclonal antibody comprises:

CDR1, CDR2 and CDR3 of a heavy chain (H chain) variable region (V region) having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively; and

CDR1, CDR2 and CDR3 of a light chain (L chain) V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

10. (Amended) The derivative of a human chimeric antibody according to claim 8, wherein the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55.

11. (Amended) The derivative of a human chimeric antibody according to claim 8, wherein the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

HANAI et al
Serial No. **Unknown**

12. (Amended) The derivative of a human chimeric antibody according to claim 8, wherein

the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55; and

the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

13. (Amended) The derivative of a human chimeric antibody KM871 according to claim 8, wherein

the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55; and

the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

17. (Amended) The derivative of a human CDR-grafted antibody according to claim 14, wherein the antibody comprises CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

18. (Amended) The derivative of a human CDR-grafted antibody according to claim 14, wherein the antibody comprises CDR1, CDR2 and CDR3 of the L chain V

HANAI et al
Serial No. **Unknown**

region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

19. (Amended) The derivative of a human CDR-grafted antibody according to claim 14, wherein the antibody comprises:

CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively; and

CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

20. (Amended) The derivative of a human CDR-grafted antibody according to claim 14, wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9.

21. (Amended) The derivative of a human CDR-grafted antibody according to claim 14, wherein the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

22. (Amended) The derivative of a human CDR-grafted antibody according to claim 14, wherein the H chain V region and the L chain V region of the antibody comprises the amino acid sequences represented by SEQ ID NO:9 and SEQ ID NO:54, respectively.

HANAI et al
Serial No. **Unknown**

23. (Amended) The derivative of a human CDR-grafted antibody KM8871 according to claim 14, wherein

the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and

the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

25. (Amended) The derivative of the antibody fragment according to claim 1, wherein the antibody fragment comprises amino acid sequences of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3 produced by a hybridoma.

26. (Amended) The derivative of the antibody fragment according to claim 1, wherein the antibody fragment comprises an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:55.

27. (Amended) The derivative of the antibody fragment according to claim 1, wherein the antibody fragment comprises an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:56.

28. (Amended) The derivative of the antibody fragment according to claim 1, wherein the antibody fragment comprises:

HANAI et al
Serial No. **Unknown**

an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:55; and

an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:56.

29. (Amended) The derivative of the antibody fragment according to claim 1, wherein the antibody fragment comprises amino acid sequences of an H chain V region and an L chain V region of a human CDR-grafted antibody against ganglioside GD3.

30. (Amended) The derivative of the antibody fragment according to claim 1, wherein the antibody fragment comprises an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:9.

31. (Amended) The derivative of the antibody fragment according to claim 1, wherein the antibody fragment comprises an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:54.

32. (Amended) The derivative of the antibody fragment according to claim 1, wherein the antibody fragment comprises:

an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:9; and

an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:54.

33. (Amended) The derivative of an antibody fragment according to claim 1, wherein the antibody fragment comprises CDR1, CDR2 and CDR3 of an H chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

34. (Amended) The derivative of the antibody fragment according to claim 1, wherein the antibody fragment comprises CDR1, CDR2 and CDR3 of an L chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

35. (Amended) The derivative of the antibody fragment according to claim 1, wherein the antibody fragment comprises:

CDR1, CDR2 and CDR3 of an H chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5; and

CDR1, CDR2 and CDR3 of an L chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8.

36. (Amended) The derivative of a monoclonal antibody or the antibody fragment thereof according to claim 1, wherein the protein is a cytokine.

HANAI et al
Serial No. **Unknown**

42. (Amended) A DNA which encodes the derivative of a monoclonal antibody or the derivative of the antibody fragment thereof which specifically reacts with ganglioside GD3 according to claim 1.

47. (Amended) A process for producing an antibody, which comprises:
culturing the transformant according to claim 44 in a culture medium to produce and accumulate the derivative of a monoclonal antibody or the derivative of the antibody fragment thereof in the culture; and
recovering the derivative of the antibody or the derivative of the antibody fragment thereof from the culture.

52. (Amended) The human CDR-grafted antibody or the antibody fragment thereof according to claim 49, wherein the antibody comprises CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

53. (Amended) The human CDR-grafted antibody or the antibody fragment thereof according to claim 49, wherein the antibody comprises CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

54. (Amended) The human CDR-grafted antibody or the antibody fragment thereof according to claim 49, wherein the antibody comprises:

HANAI et al
Serial No. **Unknown**

CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively; and

CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8.

55. (Amended) The human CDR-grafted antibody or the antibody fragment thereof according to claim 49, wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9.

56. (Amended) The human CDR-grafted antibody or the antibody fragment thereof according to claim 49, wherein the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

57. (Amended) The human CDR-grafted antibody or the antibody fragment thereof according to claim 49, wherein

the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and

the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

58. (Amended) The human CDR-grafted antibody KM8871 or the antibody fragment thereof according to claim 49, wherein

HANAI et al
Serial No. **Unknown**

the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and

the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

59. (Amended) A DNA which encodes the human CDR-grafted antibody or the antibody fragment thereof which specifically reacts with ganglioside GD3 according to claim 48.

63. (Amended) A process for producing an antibody, which comprises:
culturing the transformant according to claim 61 in a culture medium to produce and accumulate the human CDR-grafted antibody or the antibody fragment thereof in the culture; and
recovering the antibody or the antibody fragment thereof from the culture.

64. (Amended) A medicament comprising at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to claim 1 and the human CDR-grafted antibody and the antibody fragment thereof which specifically reacts with ganglioside GD3.

65. (Amended) A therapeutic agent for cancers, comprising, as an active ingredient, at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to claim 1 and the human CDR-

HANAI et al
Serial No. **Unknown**

grafted antibody and the antibody fragment thereof which specifically reacts with ganglioside GD3.

66. (Amended) A diagnostic agent for cancers, comprising, as an active ingredient, at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to claim 1 and the human CDR-grafted antibody and the antibody fragment thereof which specifically reacts with ganglioside GD3.

HANAI et al
Serial No. **Unknown**

REMARKS


Attached hereto is a marked-up version of the changes made to the claims by the current amendment. The attached page is captioned "**Version with markings to show changes made.**"

The above amendments are made to place the claims in a more traditional format.

Respectfully submitted,

NIXON & VANDERHYE P.C.

By: _____



B. J. Sadoff

Reg. No. **36,663**

BJS:Imy

1100 North Glebe Road, 8th Floor
Arlington, VA 22201-4714
Telephone: (703) 816-4000
Facsimile: (703) 816-4100

VERSION WITH MARKINGS TO SHOW CHANGES MADE

3. (Amended) The derivative of an antibody according to claim 1 [or 2], wherein the monoclonal antibody comprises CDR1, CDR2 and CDR3 of H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

4. (Amended) The derivative of an antibody according to claim 1 [or 2], wherein the monoclonal antibody comprises CDR1, CDR2 and CDR3 of L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

5. (Amended) The derivative of an antibody according to claim 1 [or 2], wherein the monoclonal antibody comprises:

CDR1, CDR2 and CDR3 of a heavy chain (H chain) variable region (V region) having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively; and

CDR1, CDR2 and CDR3 of a light chain (L chain) V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

10. (Amended) The derivative of a human chimeric antibody according to claim 8 [or 9], wherein the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55.

HANAI et al
Serial No. **Unknown**

11. (Amended) The derivative of a human chimeric antibody according to claim 8 [or 9], wherein the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

12. (Amended) The derivative of a human chimeric antibody according to claim 8 [or 9], wherein
the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55; and
the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

13. (Amended) The derivative of a human chimeric antibody KM871 according to claim 8 [or 9], wherein
the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55; and
the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

17. (Amended) The derivative of a human CDR-grafted antibody according to [any one of claims 14 to 16] claim 14, wherein the antibody comprises CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

18. (Amended) The derivative of a human CDR-grafted antibody according to [any one of claims 14 to 16] claim 14, wherein the antibody comprises CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

19. (Amended) The derivative of a human CDR-grafted antibody according to [any one of claims 14 to 16] claim 14, wherein the antibody comprises:

CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively; and

CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

20. (Amended) The derivative of a human CDR-grafted antibody according to [any one of claims 14 to 16] claim 14, wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9.

21. (Amended) The derivative of a human CDR-grafted antibody according to [any one of claims 14 to 16] claim 14, wherein the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

22. (Amended) The derivative of a human CDR-grafted antibody according to [any one of claims 14 to 16] claim 14, wherein the H chain V region and the L chain V

HANAI et al
Serial No. **Unknown**

region of the antibody comprises the amino acid sequences represented by SEQ ID NO:9 and SEQ ID NO:54, respectively.

23. (Amended) The derivative of a human CDR-grafted antibody KM8871 according to [any one of claims 14 to 16] claim 14, wherein

the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and

the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

25. (Amended) The derivative of the antibody fragment according to claim 1 [or 24], wherein the antibody fragment comprises amino acid sequences of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3 produced by a hybridoma.

26. (Amended) The derivative of the antibody fragment according to claim 1 [or 24], wherein the antibody fragment comprises an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:55.

27. (Amended) The derivative of the antibody fragment according to claim 1 [or 24], wherein the antibody fragment comprises an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:56.

HANAI et al
Serial No. **Unknown**

28. (Amended) The derivative of the antibody fragment according to claim 1 [or 24], wherein the antibody fragment comprises:

an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:55; and

an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:56.

29. (Amended) The derivative of the antibody fragment according to claim 1 [or 24], wherein the antibody fragment comprises amino acid sequences of an H chain V region and an L chain V region of a human CDR-grafted antibody against ganglioside GD3.

30. (Amended) The derivative of the antibody fragment according to claim 1 [or 24], wherein the antibody fragment comprises an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:9.

31. (Amended) The derivative of the antibody fragment according to claim 1 [or 24], wherein the antibody fragment comprises an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:54.

32. (Amended) The derivative of the antibody fragment according to claim 1 [or 24], wherein the antibody fragment comprises:

HANAI et al
Serial No. **Unknown**

an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:9; and

an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:54.

33. (Amended) The derivative of an antibody fragment according to claim 1 [or 24], wherein the antibody fragment comprises CDR1, CDR2 and CDR3 of an H chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

34. (Amended) The derivative of the antibody fragment according to claim 1 [or 24], wherein the antibody fragment comprises CDR1, CDR2 and CDR3 of an L chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

35. (Amended) The derivative of the antibody fragment according to claim 1 [or 24], wherein the antibody fragment comprises:

CDR1, CDR2 and CDR3 of an H chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5; and

CDR1, CDR2 and CDR3 of an L chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8.

36. (Amended) The derivative of a monoclonal antibody or the antibody fragment thereof according to [any one of claim 1 to 35] claim 1, wherein the protein is a cytokine.

42. (Amended) A DNA which encodes the derivative of a monoclonal antibody or the derivative of the antibody fragment thereof which specifically reacts with ganglioside GD3 according to [any one of claims 1 to 41] claim 1.

47. (Amended) A process for producing an antibody, which comprises:
culturing the transformant according to [any one of claims 44 to 46] claim 44 in a culture medium to produce and accumulate the derivative of a monoclonal antibody or the derivative of the antibody fragment thereof [according to any one of claims 1 to 41] in the culture; and

recovering the derivative of the antibody or the derivative of the antibody fragment thereof from the culture.

52. (Amended) The human CDR-grafted antibody or the antibody fragment thereof according to [any one of claims 49 to 51] claim 49, wherein the antibody comprises CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

53. (Amended) The human CDR-grafted antibody or the antibody fragment thereof according to [any one of claims 49 to 51] claim 49, wherein the antibody

HANAI et al
Serial No. **Unknown**

comprises CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

54. (Amended) The human CDR-grafted antibody or the antibody fragment thereof according to [any one of claims 49 to 51] claim 49, wherein the antibody comprises:

CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively; and

CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8.

55. (Amended) The human CDR-grafted antibody or the antibody fragment thereof according to [any one of claims 49 to 51] claim 49, wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9.

56. (Amended) The human CDR-grafted antibody or the antibody fragment thereof according to [any one of claims 49 to 51] claim 49, wherein the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

57. (Amended) The human CDR-grafted antibody or the antibody fragment thereof according to [any one of claims 49 to 51] claim 49, wherein

the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and

HANAI et al
Serial No. **Unknown**

the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

58. (Amended) The human CDR-grafted antibody KM8871 or the antibody fragment thereof according to [any one of claims 49 to 51] claim 49, wherein

the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and

the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

59. (Amended) A DNA which encodes the human CDR-grafted antibody or the antibody fragment thereof which specifically reacts with ganglioside GD3 according to [any one of claims 48 to 58] claim 48.

63. (Amended) A process for producing an antibody, which comprises:
culturing the transformant according to claim 61 [or 62] in a culture medium to produce and accumulate the human CDR-grafted antibody or the antibody fragment thereof [according to any one of claims 48 to 58] in the culture; and
recovering the antibody or the antibody fragment thereof from the culture.

64. (Amended) A medicament comprising at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to [claims 1 to 41] claim 1 and the human CDR-grafted antibody and the

HANAI et al
Serial No. **Unknown**

antibody fragment thereof [according to any one of claims 48 to 58] which specifically reacts with ganglioside GD3.

65. (Amended) A therapeutic agent for cancers, comprising, as an active ingredient, at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to [any one of claims 1 to 41] claim 1 and the human CDR-grafted antibody and the antibody fragment thereof [according to any one of claims 48 to 58] which specifically reacts with ganglioside GD3.

66. (Amended) A diagnostic agent for cancers, comprising, as an active ingredient, at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to [any one of claims 1 to 41] claim 1 and the human CDR-grafted antibody and the antibody fragment thereof [according to any one of claims 48 to 58] which specifically reacts with ganglioside GD3.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of

HANAI et al

Atty. Ref.: **249-255**

Serial No. **Unknown**

Group:

National Phase of: **PCT/JP00/06774**

International Filing Date: **29 September 2000**

Filed: **March 29, 2002**

Examiner:

For: **COMPLEMENTARY DETERMINING REGION-GRAFTED
ANTIBODY AGAINST GANGLIOSIDE GD3 AND
DERIVATIVE OF ANTIBODY AGAINST GANGLIOSIDE
GD3**

* * * * *

March 29, 2002

Assistant Commissioner for Patents
Washington, DC 20231


Sir:

STATEMENT

The attached paper and computer readable copies of the Sequence Listing are the same. No new matter has been added.

Respectfully submitted,

NIXON & VANDERHYE P.C.

By: 

B. J. Sadoff

Reg. No. **36,663**

1100 North Glebe Road, 8th Floor
Arlington, VA 22201-4714
Telephone: (703) 816-4000
Facsimile: (703) 816-4100

10/089500
JC13 Rec'd PCT/PTO 29 MAR 2002

41/p1b

SPECIFICATION

COMPLEMENTARITY DETERMINING REGION-GRAFTED ANTIBODY AGAINST GANGLIOSIDE GD3 AND DERIVATIVE OF ANTIBODY AGAINST GANGLIOSIDE GD3

TECHNICAL FIELD

The present invention relates to derivatives of a monoclonal antibody and the antibody fragment which specifically react with ganglioside GD3 (hereinafter referred to as "GD3"), and to a human CDR-grafted antibody and the antibody fragment which specifically react with GD3. Furthermore, the present invention relates to a DNA sequence which encodes the derivatives, antibody or antibody fragment. The present invention relates to a vector comprising the DNA sequence and a cell transformed with the vector. Moreover, the present invention relates to a process for producing the derivatives, antibody and antibody fragment using the transformed cell, and to a diagnostic agent and a therapeutic agent for cancers using the derivatives, antibody and antibody fragment.

BACKGROUND OF THE INVENTION

It is known that, when an antibody derived from a non-human animal, e.g., a mouse antibody, is administered to human, it is recognized as a foreign substance and thus induces a human antibody against the mouse antibody in the human body (human anti mouse antibody, hereinafter referred to as "HAMA"). It is known that the HAMA reacts with the administered mouse antibody to cause side effects (*J. Clin. Oncol.*, 2, 881 (1984); *Blood*, 65, 1349 (1985); *J. Natl. Cancer Inst.*, 80, 932 (1988); *Proc. Natl. Acad. Sci. U.S.A.*, 82, 1242 (1985)), to quicken disappearance of the administered mouse antibody from blood (*J. Nucl. Med.*, 26, 1011 (1985); *Blood*, 65, 1349 (1985); *J. Natl. Cancer Inst.*,

20, 937 (1988)) and to reduce therapeutic effects of the mouse antibody (*J. Immunol.*, 135, 1530 (1985); *Cancer Res.*, 46, 6489 (1986)).

In order to resolve these problems, attempts have been made to genetic engineeringly convert an antibody derived from a non-human animal into a humanized antibody such as a human chimeric antibody or a human complementarity determining region (hereinafter referred to as "CDR")-grafted antibody. The human chimeric antibody is an antibody in which its antibody variable region (hereinafter referred to as "V region") is derived from a non-human animal antibody and its constant region (hereinafter referred to as "C region") is derived from a human antibody (*Proc. Natl. Acad. Sci. U.S.A.*, 81, 6851 (1984)). The human CDR-grafted antibody is an antibody in which the amino acid sequence of CDR in the V region derived from a non-human animal antibody is grafted to an appropriate position of a human antibody (*Nature*, 321, 522 (1986)). These humanized antibodies have various advantages in clinically applying to human, in comparison with antibodies of non-human animals such as mouse antibodies and the like. For example, it has been reported on the immunogenicity and stability in blood that, when a human chimeric antibody is administered to human, its half-life in blood increased about six times in comparison with a mouse antibody (*Proc. Natl. Acad. Sci. U.S.A.*, 86, 4220 (1989)). It has been reported that, in a test of a human CDR-grafted antibody in monkeys, its immunogenicity is reduced and its half-life in blood is increased four to five times in comparison with a mouse antibody (*J. Immunol.*, 147, 1352 (1991)). That is, it is expected that humanized antibodies have less side effects and their therapeutic effects continue for a longer period of time than antibodies of non-human animals. In addition, when applications particularly as anti-tumor antibodies are

taken into consideration, higher cytotoxic activities such as complement-dependent cytotoxicity (hereinafter referred to as "CDC"), antibody-dependent cell-mediated cytotoxic activity (hereinafter referred to as "ADCC") and the like via an antibody Fc region (region in and after the hinge region of an antibody heavy chain) are important for the therapeutic effects. It has been reported that on such cytotoxic activities, that the Fc region of human antibodies is more superior in human to the Fc region of antibodies of non-human animals, since the Fc region more effectively activates human complement components and human effector cells having Fc receptors of mononuclear cells, macrophages and NK cells on the cell surface. For example, it has been reported that when a mouse antibody against GD3 (hereinafter referred to as "anti-GD3 mouse antibody") is converted into a human chimeric antibody having a human antibody Fc region (hereinafter referred to as "anti-GD3 chimeric antibody"), its tumor cell growth-inhibitory activity by human effector cells is increased (*J. Immunol.*, 144, 1382 (1990)), and similar results have been reported on a human CDR-grafted antibody against CAMPATH-1 antigen (*Nature*, 332, 323 (1988)).

These results clearly show that humanized antibodies are preferred to non-human antibodies such as mouse antibodies and the like, as antibodies for clinical applications to human.

In addition, with the recent advance in protein engineering and genetic engineering, antibody fragments having a smaller molecular weight such as Fab, Fab', F(ab')₂, a single chain antibody (*Science*, 242, 423 (1988)), a disulfide stabilized V region fragment (*Molecular Immunol.*, 32, 249 (1995)) and the like, can be produced. Since these fragments have a smaller molecular weight than whole antibody molecules, they are excellent in penetrating ability into target tissues (*Cancer Res.*, 52, 3402 (1992)).

It is considered that these fragments derived from humanized antibodies are more desirable than those derived from non-human animal antibodies such as mouse antibodies, when used in clinical applications to human.

5 Ganglioside which is one of glycolipids containing sialic acid is a molecule which constitutes animal cell membrane and is constituted by a sugar chain as a hydrophilic side chain and a sphingosine and fatty acid as hydrophobic side chains. It is known that kinds and
10 expression levels of ganglioside vary depending on the cell types, organ species, animal species and the like. It is also known that the expression of ganglioside changes quantitatively and qualitatively in the process of malignant transformation of cells (*Cancer Res.*, 45, 2405
15 (1985)). Particularly, the GD3 relating to the present invention is present in an extremely small amount in normal cells but in a large amount in cancer cells such as melanoma, sarcoma, glioma, neuroblastoma and the like (*Proc. Natl. Acad. Sci. U.S.A.*, 77, 6114 (1980); *J. Exp. Med.*, 155,
20 1133 (1982); *Cancer Res.*, 45, 4401 (1985)), and an antibody against GD3 (hereinafter referred to as "anti-GD3 antibody") is considered to be useful for the treatment of these cancers (*Melanoma Research*, 7, S115 (1997)). Tests on its administration to melanoma patients have already
25 been carried out using R24 as an anti-GD3 mouse antibody, and its clinical effect was observed in a part of the patients but not to an expected degree because of considerably quick disappearance of the antibody from blood due to induction of HAMA (*Melanoma Research*, 7, S115
30 (1997)). Under such circumstances, in order to achieve more effective clinical applications of an anti-GD3 antibody to human, the present inventors have attempted to produce an anti-GD3 chimeric antibody having similar activity to that of a mouse antibody based on an anti-GD3
35 mouse antibody KM641 produced by the inventors, and have

succeeded in producing an anti-GD3 chimeric antibody KM871
(Japanese Published Unexamined Patent Application
No. 304989/93). It is expected that, in comparison with
the anti-GD3 mouse antibody, the anti-GD3 chimeric antibody
5 KM871 reduces immunogenicity in human, prolongs half-life
in blood and has a stronger anti-tumor effect. However, it
was found as a result of clinical tests on other anti-GD3
chimeric antibodies in human that allergy reactions are
induced due to induction of a human antibody against a V
10 region derived from a mouse antibody in the anti-GD3
chimeric antibodies (*Cancer J. From Scientific American*, 3,
S121 (1997)). Even in the anti-GD3 chimeric antibody KM871
prepared by the inventors, it is necessary to await results
of clinical tests in human to know if similar reactions are
15 induced or not, and in order to reduce a possibility of
generating such a problem as low as possible, it is
desirable to produce a human CDR-grafted antibody against
GD3 (hereinafter referred to as "anti-GD3 CDR-grafted
antibody") which is considered to have lower immunogenicity
20 than those of anti-GD3 chimeric antibodies in human.
However, to date, there are no reports on the production of
an anti-GD3 CDR-grafted antibody.

With regard to antibody fragments, a single chain
antibody derived from an anti-GD2 mouse antibody has been
25 produced (*Hybridoma*, 16, 335 (1997)), but there are no
reports on the production of other antibody fragments.
With regard to antibody fragments derived from an anti-GD3
antibody, there are no reports on the production of
antibody fragments derived from a mouse antibody and a
30 humanized antibody. Thus, if an antibody fragment derived
from a humanized antibody against GD3 (hereinafter referred
to as "anti-GD3 humanized antibody") can be produced, it is
expected that it has excellent target tissue
penetratability and the immunogenicity is reduced.

As has been described above, it is expected that a humanized antibody and the antibody fragment have effect on diagnostic and therapeutic methods by single use, but further improvement of the effects has been examined by using them in combination with other molecules. For example, cytokine is used as one of these molecules. Cytokine is a general term for various aqueous factors which control intercellular mutual reactions in immune interactions. As cytotoxic activities of antibodies such as CDC activity, ADCC activity and the like are known and the ADCC activity is effected by effector cells having Fc receptors on the cell surface such as mononuclear cells, macrophages, NK cells and the like (*J. Immunol.*, 138, 1992 (1987)). Since various kinds of cytokine activate these effector cells, their administration in combination with antibodies has been carried out to improve ADCC activity and the like of antibodies. For example, with regard to an anti-GD2 mouse antibody 14.G2a and an anti-GD2 chimeric antibody ch14.18, their administration to human in combination with a cytokine human interleukin-2 (hereinafter referred to as "hIL-2") or a human granulocyte macrophage colony-stimulating factor (hereinafter referred to as "hGM-CSF") has been carried out (*Cancer*, 80, 317 (1997); *Cancer J. From Scientific American*, 3, S121 (1997)). Regarding an anti-GD3 mouse antibody R24, combination therapy with various kinds of cytokine has also been carried out (*Cancer Res.*, 50, 7490 (1990), *Proc. Am. Soc. Clin. Oncol.*, 1186, 345 (1992), *J. Biol. Response Mod.*, 2, 319 (1990), *Proc. Am. Soc. Clin. Oncol.*, 1182, 344 (1992), *Proc. Am. Soc. Clin. Oncol.*, 1188, 346 (1992)). However, the effects as to be expected are not found in these combination therapies due to immunogenicity of mouse antibodies or side effects derived from cytokine. Thus, derivatives has been produced by conjugating an antibody or the antibody fragment thereof with a radioisotope, a

protein (cytokine, toxin, enzyme or the like), a low molecular weight agent and the like, chemically or genetic engineeringly, and their clinical applications have been examined (*New Eng. J. Med.*, 329, 459 (1993), *Anticancer Res.*, 17, 1735 (1997), *Blood*, 78, 1173 (1991), *J. Clin. Oncol.*, 15, 723 (1997), *Bioconjugate Chem.*, 1, 606 (1997), *Cancer*, 61, 881 (1988), *Jpn. J. Cancer Res.*, 85, 167 (1994), *Antibody Immunoconjugates and Radiopharmaceuticals*, 3, 60 (1990), *Surgery*, 106, 533 (1989)). It is expected that these derivatives can accumulate a radioisotope, a protein (cytokine, toxin, enzyme or the like), a low molecular weight agent or the like in the periphery of a target tissue according to the binding specificity of antibodies so that more effective diagnoses and treatments having less side effects can be obtained. For example, with regard to hIL-2 which showed a certain degree of anti-tumor effect among cytokines used in the above combination therapy, its fusion protein with an anti-GD2 chimeric antibody ch14.18 was produced genetic engineeringly, and it has been reported that its antitumor effects in tests using mice were superior to those of simultaneous administration of anti-GD2 chimeric antibody ch14.18 and hIL-2 (*Proc. Natl. Acad. Sci. U.S.A.*, 89, 1428 (1992), *Proc. Natl. Acad. Sci. U.S.A.*, 91, 9626 (1994), *Cancer Immunol. Immunother.*, 42, 88 (1996), *Blood*, 91, 1706 (1998)). However, with regard to an anti-GD3 antibody, there are no reports on the production of derivatives by conjugating it with a protein (cytokine, toxin, enzyme or the like), a radioisotope, a low molecular weight agent and the like. Accordingly, also in the case of the anti-GD3 humanized antibody and the antibody fragment of the present invention, if derivatives conjugated with a radioisotope, a protein (toxin, enzyme or the like), a low molecular weight agent and the like, including fusion proteins with various kinds of cytokine, can be produced, reduction of immunogenicity, less side

effects and more potent antitumor effects in the tumor region can be expected when administered to the human body.

DISCLOSURE OF THE INVENTION

5 In the present invention, an amino acid sequence of each CDR was identified from the amino acid sequences of H chain and L chain V regions of the anti-GD3 mouse antibody KM641 (Japanese Published Unexamined Patent Application No. 304989/93), cDNAs encoding the H chain and L chain V regions comprising the amino acid sequences of the CDRs of 10 KM641 and amino acid sequences of the framework regions (hereinafter referred to as "FR") of human antibody H chain and L chain V regions were cloned into an expression vector for animal cell having cDNA encoding human antibody H chain and L chain C regions, and then the thus constructed anti-GD3 CDR-grafted antibody expression vector was introduced 15 into an animal cell to produce a transformant KM8871 (FERM BP-6790) which produces an anti-GD3 CDR-grafted antibody KM8871 in a culture supernatant. Next, the anti-GD3 CDR-grafted antibody KM8871 was purified from a culture supernatant of the transformant KM8871 to find that the anti-GD3 CDR-grafted antibody KM8871 shows an antigen binding activity, an antigen binding specificity and a strong cytotoxic activity against human cancer cell lines, 20 which are similar to those of the anti-GD3 chimeric antibody KM871.

Furthermore, a cDNA was constructed by linking other cDNA encoding hIL-2 to the 3'-terminal of cDNA encoding the H chain of an anti-GD3 antibody, particularly 30 anti-GD3 CDR-grafted antibody KM8871, and then the cDNA and a cDNA encoding the L chain of KM8871 were cloned into an expression vector for animal cell to construct an expression vector of a fusion protein of anti-GD3 CDR-grafted antibody KM8871 with hIL-2 (hereinafter referred to as "KM8871-hIL-2"). A transformant KM8871hIL2 (FERM BP- 35

6790) which produces KM8871-hIL-2 in culture supernatant was produced by introducing the KM8871-hIL-2 expression vector into an animal cell. Furthermore, KM8871-hIL-2 was purified from a culture supernatant of the transformant KM8871hIL2 to find that KM8871-hIL-2 shows an antigen binding activity and an antigen binding specificity, which are similar to those of anti-GD3 CDR-grafted antibody KM8871, and a growth supporting activity against cell lines showing hIL-2-dependent growth, similar to that of hIL-2. Thereafter, the present invention was accomplished by finding that an activity of KM8871-hIL-2 in terms of the cytotoxic activity measured using a human peripheral blood mononuclear cell fraction is improved in comparison with that of anti-GD3 CDR-grafted antibody KM8871.

The present invention relates to the following (1) to (66).

(1) A derivative of an antibody, comprising a monoclonal antibody or the antibody fragment thereof which specifically reacts with ganglioside GD3 which is conjugated with a radioisotope, a protein or a low molecular weight agent.

(2) The derivative of an antibody according to (1), wherein the monoclonal antibody which specifically reacts with ganglioside GD3 is an antibody selected from an antibody produced by a hybridoma, a humanized antibody and a human antibody.

(3) The derivative of an antibody according to (1) or (2), wherein the monoclonal antibody comprises CDR1, CDR2 and CDR3 of H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

(4) The derivative of an antibody according to (1) or (2), wherein the monoclonal antibody comprises CDR1, CDR2 and CDR3 of L chain V region having the amino acid

sequences represented by SEQ ID NOS:6, 7 and 8, respectively.

(5) The derivative of an antibody according to (1) or (2), wherein the monoclonal antibody comprises:

5 CDR1, CDR2 and CDR3 of a heavy chain (H chain) variable region (V region) having the amino acid sequences represented by SEQ ID NOS:3, 4 and 5, respectively; and

10 CDR1, CDR2 and CDR3 of a light chain (L chain) V region having the amino acid sequences represented by SEQ ID NOS:6, 7 and 8, respectively.

(6) The derivative of an antibody according to (2), wherein the antibody produced by a hybridoma is KM641 (FERM BP-3116).

15 (7) The derivative of an antibody according to (2), wherein the humanized antibody is a human chimeric antibody or a human CDR-grafted antibody.

(8) The derivative of a human chimeric antibody according to (7), wherein the human chimeric antibody comprises an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3 produced by a hybridoma.

(9) The derivative of a human chimeric antibody according to (7), wherein the human chimeric antibody comprises:

25 an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3 produced by a hybridoma; and

an H chain constant region (C region) and an L chain C region of a human antibody.

30 (10) The derivative of a human chimeric antibody according to (8) or (9), wherein the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55.

35 (11) The derivative of a human chimeric antibody according to (8) or (9), wherein the L chain V region

comprises the amino acid sequence represented by SEQ ID NO:56.

(12) The derivative of a human chimeric antibody according to (8) or (9), wherein

5 the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55; and

the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

10 (13) The derivative of a human chimeric antibody KM871 according to (8) or (9), wherein

the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55; and

the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

15 (14) The derivative of a human CDR-grafted antibody according to (7), wherein the human CDR-grafted antibody comprises CDR of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3.

20 (15) The derivative of a human CDR-grafted antibody according to (7), wherein the human CDR-grafted antibody comprises:

CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3; and

25 framework regions (FRs) of an H chain V region and an L chain V region of a human antibody.

(16) The derivative of an antibody according to (7), wherein the human CDR-grafted antibody comprises:

30 CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3;

FRs of an H chain V region and an L chain V region of a human antibody; and

an H chain C region and an L chain C region of a human antibody.

(17) The derivative of a human CDR-grafted antibody according to any one of (14) to (16), wherein the antibody comprises CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

(18) The derivative of a human CDR-grafted antibody according to any one of (14) to (16), wherein the antibody comprises CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

(19) The derivative of a human CDR-grafted antibody according to any one of (14) to (16), wherein the antibody comprises:

CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively; and

CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

(20) The derivative of a human CDR-grafted antibody according to any one of (14) to (16), wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9.

(21) The derivative of a human CDR-grafted antibody according to any one of (14) to (16), wherein the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

(22) The derivative of a human CDR-grafted antibody according to any one of (14) to (16), wherein the H chain V region and the L chain V region of the antibody comprises the amino acid sequences represented by SEQ ID NO:9 and SEQ ID NO:54, respectively.

(23) The derivative of a human CDR-grafted antibody KM8871 according to any one of (14) to (16), wherein

the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and

the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

5 (24) The derivative of the antibody fragment according to (1), wherein the antibody fragment is an antibody fragment selected from Fab, Fab', F(ab')₂, a single chain antibody (scFv), a disulfide stabilized V region fragment (dsFv) and a peptide comprising CDR.

10 (25) The derivative of the antibody fragment according to (1) or (24), wherein the antibody fragment comprises amino acid sequences of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3 produced by a hybridoma.

15 (26) The derivative of the antibody fragment according to (1) or (24), wherein the antibody fragment comprises an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:55.

20 (27) The derivative of the antibody fragment according to (1) or (24), wherein the antibody fragment comprises an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:56.

25 (28) The derivative of the antibody fragment according to (1) or (24), wherein the antibody fragment comprises:

an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:55; and

an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:56.

30 (29) The derivative of the antibody fragment according to (1) or (24), wherein the antibody fragment comprises amino acid sequences of an H chain V region and an L chain V region of a human CDR-grafted antibody against ganglioside GD3.

(30) The derivative of the antibody fragment according to (1) or (24), wherein the antibody fragment comprises an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:9.

5 (31) The derivative of the antibody fragment according to (1) or (24), wherein the antibody fragment comprises an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:54.

10 (32) The derivative of the antibody fragment according to (1) or (24), wherein the antibody fragment comprises:

an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:9; and

15 an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:54.

20 (33) The derivative of an antibody fragment according to (1) or (24), wherein the antibody fragment comprises CDR1, CDR2 and CDR3 of an H chain V region of the antibody having the amino acid sequences represented by SEQ ID NOS:3, 4 and 5, respectively.

25 (34) The derivative of the antibody fragment according to (1) or (24), wherein the antibody fragment comprises CDR1, CDR2 and CDR3 of an L chain V region of the antibody having the amino acid sequences represented by SEQ ID NOS:6, 7 and 8, respectively.

(35) The derivative of the antibody fragment according to (1) or (24), wherein the antibody fragment comprises:

30 CDR1, CDR2 and CDR3 of an H chain V region of the antibody having the amino acid sequences represented by SEQ ID NOS:3, 4 and 5; and

CDR1, CDR2 and CDR3 of an L chain V region of the antibody having the amino acid sequences represented by SEQ ID NOS:6, 7 and 8.

(36) The derivative of a monoclonal antibody or the antibody fragment thereof according to any one of (1) to (35), wherein the protein is a cytokine.

5 (37) The derivative of a monoclonal antibody or the antibody fragment thereof according to (36), wherein the cytokine is human interleukin-2 (hIL-2).

(38) The derivative of an antibody according to (37), wherein the derivative of an antibody comprises a human chimeric antibody KM871 and hIL-2.

10 (39) The derivative of an antibody according to (38), wherein the antibody conjugated with hIL-2 comprises:

an H chain V region having the amino acid sequence represented by SEQ ID NO:57; and

15 an L chain V region having the amino acid sequence represented by SEQ ID NO:56.

(40) The derivative of an antibody according to (37), wherein the derivative of an antibody comprises a human CDR-grafted antibody KM8871 and hIL-2.

20 (41) The derivative of an antibody according to (1), wherein the antibody conjugated with hIL-2 comprises:

an H chain V region having the amino acid sequence represented by SEQ ID NO:53; and

an L chain V region having the amino acid sequence represented by SEQ ID NO:54.

25 (42) A DNA which encodes the derivative of a monoclonal antibody or the derivative of the antibody fragment thereof which specifically reacts with ganglioside GD3 according to any one of (1) to (41).

30 (43) A recombinant vector comprising the DNA according to (42).

(44) A transformant which is obtained by introducing the recombinant vector according to (43) into a host cell.

35 (45) A transformant KM871hIL2 (FERM BP-6918) which produces the antibody according to (38).

(46) A transformant KM8871hIL2 (FERM BP-6791) which produces the antibody according to (40).

(47) A process for producing an antibody, which comprises:

5 culturing the transformant according to any one of (44) to (46) in a culture medium to produce and accumulate the derivative of a monoclonal antibody or the derivative of the antibody fragment thereof according to any one of (1) to (41) in the culture; and

10 recovering the derivative of the antibody or the derivative of the antibody fragment thereof from the culture.

(48) A human CDR-grafted antibody or the antibody fragment thereof which specifically reacts with ganglioside
15 GD3.

(49) The human CDR-grafted antibody or the antibody fragment thereof according to (48), wherein the human CDR-grafted antibody comprises CDRs of an H chain V region and an L chain V region of a monoclonal antibody
20 against ganglioside GD3.

(50) The human CDR-grafted antibody or the antibody fragment thereof according to (48), wherein the human CDR-grafted antibody comprises:

25 CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3; and

FRS of an H chain V region and an L chain V region of a human antibody.

(51) The human CDR-grafted antibody or the antibody fragment thereof according to (48), wherein the
30 human CDR-grafted antibody comprises:

CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3;

FRS of an H chain V region and an L chain V region of a human antibody; and

an H chain C region and an L chain C region of a human antibody.

5 (52) The human CDR-grafted antibody or the antibody fragment thereof according to any one of (49) to (51), wherein the antibody comprises CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

10 (53) The human CDR-grafted antibody or the antibody fragment thereof according to any one of (49) to (51), wherein the antibody comprises CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

15 (54) The human CDR-grafted antibody or the antibody fragment thereof according to any one of (49) to (51), wherein the antibody comprises:

CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively; and

20 CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8.

25 (55) The human CDR-grafted antibody or the antibody fragment thereof according to any one of (49) to (51), wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9.

30 (56) The human CDR-grafted antibody or the antibody fragment thereof according to any one of (49) to (51), wherein the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

(57) The human CDR-grafted antibody or the antibody fragment thereof according to any one of (49) to (51), wherein

the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and

the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

5 (58) The human CDR-grafted antibody KM8871 or the antibody fragment thereof according to any one of (49) to (51), wherein

the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and

10 the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

(59) A DNA which encodes the human CDR-grafted antibody or the antibody fragment thereof which specifically reacts with ganglioside GD3 according to any
15 one of (48) to (58).

(60) A recombinant vector comprising the DNA according to (59).

(61) A transformant which is obtained by introducing the recombinant vector according to (60) into a
20 host cell.

(62) A transformant KM8871 (FERM BP-6790) which produces the human CDR-grafted antibody according to (58).

(63) A process for producing an antibody, which comprises:

25 culturing the transformant according to (61) or (62) in a culture medium to produce and accumulate the human CDR-grafted antibody or the antibody fragment thereof according to any one of (48) to (58) in the culture; and

recovering the antibody or the antibody fragment
30 thereof from the culture.

(64) A medicament comprising at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to (1) to (41) and the human CDR-grafted antibody and the

antibody fragment thereof according to any one of (48) to (58).

(65) A therapeutic agent for cancers, comprising, as an active ingredient, at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to any one of (1) to (41) and the human CDR-grafted antibody and the antibody fragment thereof according to any one of (48) to (58).

(66) A diagnostic agent for cancers, comprising, as an active ingredient, at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to any one of (1) to (41) and the human CDR-grafted antibody and the antibody fragment thereof according to any one of (48) to (58).

The present invention relates to derivatives of an antibody, comprising a monoclonal antibody which specifically reacts with ganglioside GD3 or the antibody fragment thereof which is conjugated with a radioisotope, a protein or a low molecular weight agent.

Examples of the monoclonal antibody include an antibody produced by a hybridoma, a humanized antibody, a human antibody and the like.

A hybridoma is a cell, which is obtained by subjecting a B cell prepared by immunizing a non-human mammal with an antigen to cell fusion with a myeloma cell derived from a mouse or the like which produces a desired monoclonal antibody having an antigen specificity and.

Examples of the humanized antibody include a human chimeric antibody, a human CDR-grafted antibody and the like.

A human chimeric antibody is an antibody which comprises an H chain V region (hereinafter referred to as "HV" or "VH") and an L chain V region (hereinafter referred to as "LV" or "VL") of an antibody derived from a non-human

animal, an H chain C region (hereinafter referred to as "CH") of a human antibody and an L chain C region (hereinafter referred to as "CL") of a human antibody. As the non-human animal, any animal such as mouse, rat, hamster, rabbit or the like can be used, so long as a hybridoma can be prepared therefrom.

The human chimeric antibody of the present invention can be produced by obtaining cDNAs encoding VH and VL from a hybridoma which produces a monoclonal antibody which specifically reacts with GD3, constructing a human chimeric antibody expression vector by inserting them into an expression vector for animal cell having genes encoding human antibody CH and human antibody CL, and expressing the expression vector by introducing it into an animal cell.

As the CH of a human chimeric antibody, it may be any region which belongs to human immunoglobulin (hereinafter referred to as "hIg"), but those of hIgG class are suitable and any one of subclasses belonging to hIgG class, such as hIgG1, hIgG2, hIgG3 and hIgG4, can also be used. Also, as the CL of a human chimeric antibody, it may be any region which belongs to hIg, and those of κ class or λ class can be used.

Examples of the anti-GD3 chimeric antibody include KM871 described in Japanese Published Unexamined Patent Application No. 304989/93.

A human CDR-grafted antibody is an antibody in which amino acid sequences of CDRs in VH and VL derived from a non-human animal antibody are grafted to appropriate positions of VH and VL of a human antibody.

The anti-GD3 CDR-grafted antibody of the present invention can be produced by constructing cDNAs encoding V regions in which CDR sequences of VH and VL of an antibody derived from a non-human animal which specifically reacts with GD3 are grafted to CDR sequences of VH and VL of an

optional human antibody, constructing a human CDR-grafted antibody expression vector by inserting them into an expression vector for animal cell having genes encoding human antibody CH and human antibody CL, and expressing the human CDR-grafted antibody by introducing the expression vector into an animal cell.

As the CH of the human CDR-grafted antibody of the present invention, it may be any region which belongs to hIg, but those of hIgG class are suitable and any one of subclasses belonging to hIgG class, such as hIgG1, hIgG2, hIgG3 and hIgG4, can also be used. Also, as the CL of the human CDR-grafted antibody, it may be any region which belongs to hIg, and those of κ class or λ class can be used.

Examples of the anti-GD3 CDR-grafted antibody include antibody KM8871 in which VH of the antibody comprises the amino acid sequence represented by SEQ ID NO:9 and CH thereof comprises an amino acid sequence of a human antibody IgG1 subclass, and VL of the antibody comprises the amino acid sequence represented by SEQ ID NO:54 and CL thereof comprises an amino acid sequence of a human antibody κ class.

Originally, a human antibody is an antibody naturally present in the human body. It also includes antibodies obtained from a human antibody phage library and a human antibody-producing transgenic animal which are produced based on the recent advance in genetic engineering, cell engineering and developmental engineering techniques.

With regard to the antibody present in the human body, a lymphocyte capable of producing the antibody can be cultured by isolating a human peripheral blood lymphocyte and infecting it with EB virus or the like to thereby immortalizing it, followed by cloning, and the antibody can be purified from the culture mixture.

The human antibody phage library is a library in which antibody fragments such as Fab, scFv and the like are

expressed on the phage surface by inserting an antibody gene prepared from human B cell into a phage gene. From the library, a phage expressing a desired antibody fragment having an antigen binding activity can be recovered, using
5 its activity to bind to an antigen-immobilized substrate as the marker. The antibody fragment can be genetic engineeringly converted further into a human antibody molecule having two complete H chains and two complete L chains.

10 A human antibody-producing transgenic non-human animal is an animal in which a human antibody gene is introducing into cells. Specifically, a human antibody-producing transgenic animal can be produced by introducing a human antibody gene into mouse ES cell, transplanting the
15 ES cell into an early stage embryo of other mouse and then developing it. As the preparation method of a human antibody from the human antibody-producing transgenic animal, the human antibody can be produced and accumulated in a culture mixture by obtaining a human antibody-producing hybridoma by a hybridoma production method usually carried out in non-human mammals and then culturing
20 it.

Examples of the antibody fragment include Fab, Fab', F(ab')₂, scFv, dsFv, a peptide comprising CDR and the like.

25 An Fab is an antibody fragment having a molecular weight of about 50,000 and antigen binding activity, in which about a half of the N-terminal side of H chain and the entire L chain, among fragments obtained by treating IgG with a protease, papain (cut at the 224th position amino acid residue of the H chain), are bound together
30 through a disulfide bond.

The Fab of the present invention can be obtained treating an antibody which specifically reacts with GD3 with a protease, papain. Also, the Fab can be produced by
35 inserting DNA encoding Fab of the antibody into an

expression vector for procaryote or an expression vector for eucaryote, and introducing the vector into a procaryote or eucaryote to express the Fab.

5 An $F(ab')_2$ is an antibody fragment having a molecular weight of about 100,000 and antigen binding activity, which is slightly larger than the Fab bound via a disulfide bond of the hinge region, among fragments obtained by treating IgG with a protease, pepsin (cut at the 234th position amino acid residue of the H chain).

10 The $F(ab')_2$ of the present invention can be obtained treating an antibody which specifically reacts with GD3 with a protease, pepsin. Also, the $F(ab')_2$ can be produced by binding Fab' described below via a thioether bond or a disulfide bond.

15 An Fab' is an antibody fragment having a molecular weight of about 50,000 and antigen binding activity, which is obtained by cutting a disulfide bond of the hinge region of the $F(ab')_2$.

20 The Fab' of the present invention can be obtained treating $F(ab')_2$ which specifically reacts with GD3 with a reducing agent, dithiothreitol. Also, the Fab' can be produced by inserting DNA encoding Fab' fragment of the antibody into an expression vector for procaryote or an expression vector for eucaryote, and introducing the vector
25 into a procaryote or eucaryote to effect its expression.

30 An scFv is a VH-P-VL or VL-P-VH polypeptide in which one chain VH and one chain VL are linked using an appropriate peptide linker (hereinafter referred to as "P"). The VH and VL in the scFv used in the present invention may be any antibody of an antibody produced by a hybridoma, a humanized antibody and a human antibody.

35 The scFv of the present invention can be produced by obtaining cDNA encoding the VH and VL of an antibody which specifically reacts with GD3, constructing DNA encoding scFv, inserting the DNA into an expression vector

for procaryote or an expression vector for eucaryote, and then introducing the expression vector into a prokaryote or eucaryote to express the scFv.

5 A dsFv is obtained by binding polypeptides in which one amino acid residue of each of VH and VL is substituted with a cysteine residue via a disulfide bond between the cysteine residues. The amino acid residue to be substituted with a cysteine residue can be selected based on a three-dimensional structure estimation of the antibody
10 in accordance with the method shown by Reiter et al. (*Protein Engineering*, 7: 697 (1994)). The VH and VL in the dsFv of the present invention may be any antibody of an antibody produced by a hybridoma, a humanized antibody and a human antibody.

15 The dsFv of the present invention can be produced by obtaining cDNA encoding the VH and VL of an antibody capable of specifically reacting with GD3, constructing DNA encoding dsFv, inserting the DNA into an expression vector for procaryote or an expression vector for eucaryote, and
20 then introducing the expression vector into a prokaryote or eucaryote to express the dsFv.

A peptide comprising CDR is constituted by at least one region of H chain and L chain CDRs. Plural CDRs can be bound directly or via an appropriate peptide linker.

25 The peptide comprising CDR of the present invention can be produced by obtaining cDNA encoding the VH and VL of an antibody capable of specifically reacting with GD3, inserting the cDNA into an expression vector for prokaryote or an expression vector for eucaryote, and then introducing
30 the expression vector into a prokaryote or eucaryote to express the peptide comprising CDR.

The peptide comprising CDR can also be produced by a chemical synthesis method such as Fmoc method (fluorenylmethoxycarbonyl method) or tBoc method (t-
35 butyloxycarbonyl method).

The antibody derivatives of the present invention can be produced by chemically conjugating a radioisotope, a protein, a low molecular weight agent or the like to the N-terminal side or C-terminal side of the H chain or L chain of the antibody or the antibody fragment which specifically reacts with GD3, to an appropriate substituent or side chain of the antibody or the antibody fragment, or to a sugar chain of the antibody or the antibody fragment (*Antibody Engineering Handbook*, edited by Osamu Kanemitsu, published by Chijin Shokan (1994)).

Also, the antibody derivatives can be genetically produced by linking a DNA encoding an antibody or the antibody fragment which specifically reacts with GD3 to other DNA encoding a protein to be bound, inserting the DNA into an expression vector, and introducing the expression vector into a host cell.

Examples of the isotope include ^{131}I , ^{125}I and the like, and they can be conjugated to an antibody by, e.g., a chloramine T method.

Examples of the low molecular weight agent include anticancer agents such as alkylating agent (e.g., nitrogen mustard, cyclophosphamide, etc.), metabolic antagonist (e.g., 5-fluorouracil, methotrexate, etc.), antibiotic (e.g., daunomycin, bleomycin, mitomycin C, daunorubicin, doxorubicin, etc.), plant alkaloid (e.g., vincristine, vinblastine, vindesine, etc.), hormone drug (e.g., tamoxifen, dexamethasone, etc.) and the like (*Clinical Oncology*, edited by Japanese Society of Clinical Oncology, published by Cancer and Chemotherapy (1996)); anti-inflammatory agents such as steroid agent (e.g., hydrocortisone, prednisone, etc.), non-steroidal drug (e.g., aspirin, indometacin, etc.), immunomodulator (e.g., aurothiomalate, penicillamine, etc.), immunosuppressing agent (e.g., cyclophosphamide, azathioprine, etc.), antihistaminic agent (e.g., chlorpheniramine maleate,

clemastine, etc.) and the like (*Inflammation and Anti-inflammatory Therapy*, Ishiyaku Shuppan (1982)); and the like. Examples of the method for conjugating daunomycin to an antibody include a method in which daunomycin and an amino group of an antibody are conjugated via glutaraldehyde, a method in which an amino group of daunomycin and a carboxyl group of an antibody are conjugated via a water-soluble carbodiimide, and the like.

As the protein, cytokine which activates immunocompetent cells is preferred, and examples include hIL-2, hGM-CSF, human macrophage colony-stimulating factor (hereinafter referred to as "hM-CSF"), human interleukin 12 (hereinafter referred to as "hIL-12") and the like. Also, in order to injure cancer cells directly, toxins such as ricin, diphtheria toxin and the like can be used. For example, a fusion antibody with a protein can be produced by conjugating a cDNA encoding an antibody or the antibody fragment to other cDNA encoding the protein, constructing a DNA encoding the fusion antibody, inserting the DNA into an expression vector for prokaryote or an expression vector for eucaryote, and then introducing the expression vector into a prokaryote or eucaryote to express the fusion antibody.

Examples of the fusion protein of the anti-GD3 human antibody with cytokine include a fusion protein of anti-GD3 CDR-grafted antibody KM8871 with hIL-2 and a fusion protein of anti-GD3 chimeric antibody KM871 with hIL-2. Examples of the fusion protein of anti-GD3 CDR-grafted antibody KM8871 with hIL-2 include KM8871-hIL-2 in which the H chain V region of the antibody conjugated with hIL-2 has the amino acid sequence represented by SEQ ID NO:53, the L chain V region of the antibody has the amino acid sequence represented by SEQ ID NO:54 and the L chain C region has an amino acid sequence of a human antibody κ class. Examples of the fusion protein of anti-GD3 chimeric

antibody KM871 with hIL-2 include KM871-hIL-2 in which the H chain V region of the antibody conjugated with hIL-2 has the amino acid sequence represented by SEQ ID NO:57, the L chain V region of the antibody has the amino acid sequence represented by SEQ ID NO:56 and the L chain C region has an amino acid sequence of a human antibody κ class.

The methods for producing a humanized antibody which specifically reacts with GD3 and derivatives of an antibody or the antibody fragment which specifically reacts with GD3.

1. Production of humanized antibody

(1) Construction of humanized antibody expression vector

The humanized antibody expression vector is an expression vector for animal cell to which genes encoding the CH and CL of a human antibody are inserted, and can be constructed by cloning the genes which encode the CH and CL of a human antibody into an expression vector for animal cell. The C regions of a human antibody can be any human antibody CH and CL, and examples include a C region belonging to IgG1 subclass of in an H chain of a human antibody (hereinafter referred to as "hC γ 1") and a C region belonging to κ class in an L chain of a human antibody (hereinafter referred to as "hC κ "). A chromosomal DNA comprising an exon and an intron can be used as the gene encoding CH and CL of a human antibody, and cDNA can also be used. Any expression vector for animal cell can be used, so long as a gene encoding the human antibody C region can be inserted and expressed. Examples include pAGE107 (Cytotechnology, 3, 133 (1990)), pAGE103 (J. Biochem., 101, 1307 (1987)), pHSG274 (Gene, 27, 223 (1984)), pKCR (Proc. Natl. Acad. Sci. U.S.A., 78, 1527 (1981)), pSG1 β d2-4 (Cytotechnology, 4, 173 (1990)) and the like. Examples of the promoter and the enhancer used in the expression vector for animal cell include early promoter and enhancer of SV40

(*J. Biochem.*, 101, 1307 (1987), LTR promoter and enhancer of Moloney mouse leukemia virus (*Biochem. Biophys. Res. Commun.*, 149, 960 (1987)), promoter (*Cell*, 41, 479, 1985)) and enhancer (*Cell*, 33, 717 (1983)) of immunoglobulin H chain and the like.

The humanized antibody expression vector may be either of a type in which a gene encoding an antibody H chain and a gene encoding an antibody L chain exist on separate vectors or of a type in which both genes exist on the same vector (tandem type). In respect of easiness of construction of a humanized antibody expression vector, easiness of introduction into animal cells, and balance between the expression levels of antibody H and L chains in animal cells, a tandem type of the humanized antibody expression vector is more preferable (*J. Immunol. Methods*, 167, 271 (1994)). Examples of the tandem type humanized antibody expression vector include pKANTEX93 (WO 97/10354), pEE18 (*HYBRIDOMA*, 17, 559 (1998)) and the like.

The thus constructed humanized antibody expression vector can be used for the expression of the human chimeric antibody and human CDR-grafted antibody in animal cells.

(2) Preparation of cDNA encoding V region of antibody derived from non-human animal

CDNAs encoding VH and VL of an antibody derived from a non-human animal such as a mouse antibody are obtained as follows.

mRNA is extracted from hybridoma cells producing a mouse antibody or the like to synthesize cDNA. The synthesized cDNA is inserted into a vector such as a phage, a plasmid or the like, to prepare a cDNA library. Each of a recombinant phage or recombinant plasmid containing cDNA encoding VH and a recombinant phage or recombinant plasmid containing cDNA encoding VL is isolated from the library using a C region part or a V region part of a mouse

antibody as the DNA probe. The full nucleotide sequences of the VH and VL of the mouse antibody of interest on the recombinant phage or recombinant plasmid are determined, and the full amino acid sequences of the VH and VL are deduced from the nucleotide sequences.

The non-human animal may be any animal such as mouse, rat, hamster, rabbit or the like, so long as a hybridoma cell can be produced therefrom. Examples of the method for preparing total RNA from a hybridoma cell include a guanidine thiocyanate-caesium trifluoroacetate method (*Methods in Enzymol.*, 154, 3 (1987)) and the like. Examples of the method for preparing mRNA from total RNA include an oligo (dT) immobilized cellulose column method (*Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Lab. Press, New York (1989), hereinafter referred to as "*Molecular Cloning: A Laboratory Manual*") and the like. Also, examples of a kit for preparing mRNA from a hybridoma cell include Fast Track mRNA Isolation Kit (manufactured by Invitrogen), Quick Prep mRNA Purification Kit (manufactured by Pharmacia) and the like.

Examples of the method for synthesizing cDNA and preparing a cDNA library include known methods (*Molecular Cloning: A Laboratory Manual*; *Current Protocols in Molecular Biology*, Supplement 1-34); a method using a commercially available kit such as Super Script™ Plasmid System for cDNA Synthesis and Plasmid Cloning (manufactured by GIBCO BRL), ZAP-cDNA Kit (manufactured by Stratagene), etc.; and the like.

The vector into which the cDNA synthesized using mRNA extracted from a hybridoma cell as the template is inserted for preparing a cDNA library may be any vector, so long as the cDNA can be inserted. Examples include ZAP Express (*Strategies*, 5: 58 (1992)), pBluescript II SK(+) (*Nucleic Acids Research*, 17: 9494 (1989)), λzapII (manufactured by Stratagene), λgt10 and λgt11 (*DNA Cloning*:

A Practical Approach, I, 49 (1985)), Lambda BlueMid (manufactured by Clontech), λExCell and pT7T3 18U (manufactured by Pharmacia), pcD2 (Mol. Cell. Biol., 3: 280 (1983)), pUC18 (Gene, 33: 103 (1985)), and the like.

5 Any *E. coli* for introducing the cDNA library constructed by a phage or plasmid vector may be used, so long as the cDNA library can be introduced, expressed and maintained. Examples include XL1-Blue MRF' (Strategies, 5: 81 (1992)), C600 (Genetics, 39: 440 (1954)), Y1088 and
10 Y1090 (Science, 222: 778 (1983)), NM522 (J. Mol. Biol., 166: 1 (1983)), K802 (J. Mol. Biol., 16: 118 (1966)), JM105 (Gene, 38: 275 (1985)), and the like.

For selecting cDNA clones encoding VH and VL of an antibody derived from a non-human animal in the cDNA
15 library, a colony hybridization or plaque hybridization method using an isotope- or fluorescence-labeled probe may be used (Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Lab. Press, New York (1989)). Also, the cDNAs encoding VH and VL can be prepared through polymerase
20 chain reaction (hereinafter referred to as "PCR"; Molecular Cloning: A Laboratory Manual; Current Protocols in Molecular Biology) by preparing primers and using cDNA prepared from mRNA or a cDNA library as the template.

The nucleotide sequence of the cDNA can be
25 determined by digesting the cDNA selected by the above method with appropriate restriction enzymes and the like, cloning the fragments into a plasmid such as pBluescript SK(-) (manufactured by Stratagene) or the like, carrying out the reaction by a usually used nucleotide analyzing
30 method such as the dideoxy method of Sanger, F. et al. (Proc. Natl. Acad. Sci. USA, 74: 5463 (1977)) or the like, and then analyzing the sequence using an automatic nucleotide sequence analyzer such as A.L.F. DNA sequencer (manufactured by Pharmacia) or the like.

Whether the obtained cDNAs encode the full amino acid sequences of the VH and VL of the antibody containing a secretory signal sequence can be confirmed by estimating the full amino acid sequences of the VH and VL from the determined nucleotide sequence and comparing them with full amino acid sequences of the VH and VL of known antibodies (*Sequences of Proteins of Immunological Interest*, US Dept. Health and Human Services (1991), hereinafter referred to as "*Sequences of Proteins of Immunological Interest*").

(3) Analysis of amino acid sequence of V region derived from a non-human animal

Regarding the complete amino acid sequences of VH and VL of the antibody comprising a secretory signal sequence, the length of the secretory signal sequence and N-terminal amino acid sequences can be deduced and subgroups to which they belong can also be known, by comparing with full amino acid sequences of VH and VL of known antibodies (*Sequences of Proteins of Immunological Interest*). Also, regarding the amino acid sequence of each CDR of VH and VL, it can be found by comparing it with amino acid sequences of VH and VL of known antibodies (*Sequences of Proteins of Immunological Interest*).

(4) Construction of human chimeric antibody expression vector

A human chimeric antibody expression vector can be constructed by cloning cDNAs encoding VH and VL of an antibody derived from a non-human animal in the region upstream of genes encoding CH and CL of the human antibody on the humanized antibody expression vector as described in the item 1(1). For example, each of cDNAs encoding VH and VL of an antibody derived from a non-human animal is linked with a synthesized DNA comprising nucleotide sequences at the 3' terminals of VH and VL of an antibody derived from a

non-human animal and nucleotide sequences at the 5' terminals of CH and CL of a human antibody and having a recognition sequence of an appropriate restriction enzyme at both terminals, and each cDNA is cloned so that it is appropriately expressed in upstream of genes encoding the CH and CL of the humanized antibody expression vector as described in the item 1(1) to thereby construct a human chimeric antibody expression vector.

10 (5) Construction of cDNA encoding V region of human CDR-grafted antibody

cDNAs encoding VH and VL of a human CDR-grafted antibody can be obtained as follows. First, amino acid sequences of FRs in VH and VL of a human antibody to which amino acid sequences of CDRs in VH and VL of an antibody derived from a non-human animal antibody are grafted are selected. Any amino acid sequences of FRs in VH and VL of a human antibody can be used, so long as they are derived from human. Examples include amino acid sequences of FRs in VH and VL of human antibodies registered in database such as Protein Data Bank or the like, and amino acid sequences common to subgroups of FRs in the VH and VL of human antibodies (*Sequences of Proteins of Immunological Interest*) and the like. In order to produce a human CDR-grafted antibody having potent activity, amino acid sequences having high homology (at least 60% or more) with amino acid sequence of FRs of VH and VL of an antibody of interest derived from a non-human animal is preferably selected. Then, amino acid sequences of CDRs of VH and VL of the antibody derived from a non-human animal are grafted to the selected amino acid sequences of FRs of VH and VL of a human antibody to design amino acid sequences of the VH and VL of a human CDR-grafted antibody. The designed amino acid sequences are converted to DNA sequences by considering the frequency of codon usage found in

nucleotide sequences of genes of antibodies (*Sequence of Proteins of Immunological Interest*), and the DNA sequences encoding the amino acid sequences of the VH and VL of a human CDR-grafted antibody are designed. Several synthetic
5 DNAs having a length of about 100 nucleotides are synthesized, and PCR is carried out using them. In this case, it is preferred in each of the H chain and the L chain that 6 synthetic DNAs are designed in view of the reaction efficiency of PCR and the lengths of DNAs which
10 can be synthesized.

Furthermore, they can be easily cloned into the humanized antibody expression vector constructed in the item 1(1) by introducing the recognition sequence of an appropriate restriction enzyme to the 5' terminal of the
15 synthetic DNAs present on the both terminals. After the PCR, an amplified product is cloned into a plasmid such as pBluescript SK (-) (manufactured by Stratagene) or the like, and the nucleotide sequences are determined according to the method described in the item 1(2) to obtain a plasmid
20 having DNA sequences encoding the VH and VL of a designed human CDR-grafted antibody.

(6) Modification of amino acid sequence of V region of human CDR-grafted antibody

25 It is known that when a human CDR-grafted antibody is produced by simply grafting only CDRs in VH and VL an antibody derived from a non-human animal in FRs of the VH and VL of a human antibody, its antigen binding activity is reduced in comparison with that of the original antibody
30 derived from a non-human animal (*BIO/TECHNOLOGY*, 2, 266 (1991)). As the reason, it is considered that several amino acid residues in not only CDRs but also FRs directly or indirectly relate to the antigen binding activity in the VH and VL of the original antibody derived from a non-human
35 animal, and that these amino acid residues are changed to

different amino acid residues derived from FRs of the VH and VL of the human antibody. In order to resolve the problem, in human CDR-grafted antibodies, attempts have been made to identify, among amino acid sequences of the FR of the VH and VL of human antibodies, an amino acid residue which directly relates to binding to the antibody, an amino acid residue which is interacted with an amino acid residue of CDR or an amino acid residue which keeps three-dimensional structure of the antibody and is directly related to its binding to the antigen, and to increase the reduced antigen binding activity by changing them into amino acid residues of the original antibody derived from a non-human animal (*BIO/TECHNOLOGY*, 2, 266 (1991)). In producing a human CDR-grafted antibody, it is the most important point to efficiently identify these FR amino acid residues related to the antigen binding activity, so that construction and analysis of the three-dimensional structure of antibodies have been carried out by X-ray crystallography (*J. Mol. Biol.*, 112, 535 (1977)), computer-modeling (*Protein Engineering*, 7, 1501 (1994)) and the like. Although the information of the three-dimensional structure of antibodies has been useful in the production of a human CDR-grafted antibody, no method for producing a human CDR-grafted antibody which can be applied to any antibodies has been established yet. Therefore, various attempts must be currently be necessary, for example, several modified antibodies of each antibody are produced and the relationship between each of the modified antibodies and its antibody binding activity is examined.

Substitution of the amino acid residues of the FR in the VH and VL of a human antibody can be achieved by carrying out the PCR described in the item 1(5) using synthetic DNA for modification. With regard to an amplified product obtained by the PCR, the nucleotide sequence is determined according to the method described in

the item 1(2) so that whether the objective modification has been carried out is confirmed.

5 (7) Construction of human CDR-grafted antibody expression vector

A human CDR-grafted antibody expression vector can be constructed by cloning cDNAs encoding VH and VL of the human CDR-grafted antibody constructed in the items 1(5) and 1(6) into upstream of the genes encoding CH and CL of the human antibody in the humanized antibody expression vector as described in the item 1(1).

10 For example, when recognition sites for an appropriate restriction enzymes are introduced to the 5'-terminal of synthetic DNAs positioned at both terminals among synthetic DNAs used in the construction of VH and VL of the human CDR-grafted antibody in the items 1(5) and 1(6), cloning can be carried out so that they are expressed in an appropriate form in upstream of genes encoding CH and CL of the human antibody in the humanized antibody expression vector as described in the item 1(1).

(8) Transient expression of humanized antibody

25 In order to efficiently evaluate the antigen binding activity of various humanized antibodies produced, the humanized antibodies can be expressed transiently using the humanized antibody expression vector as described in the items 1(4) and 1(7) or the modified expression vector thereof. Any cell can be used as a host cell, so long as the host cell can express a humanized antibody. Generally, 30 COS-7 cell (ATCC CRL1651) is used in view of its high expression level (*Methods in Nucleic Acids Res.*, CRC Press, 283 (1991)). Examples of the method for introducing the expression vector into COS-7 cell include a DEAE-dextran method (*Methods in Nucleic Acids Res.*, CRC Press, 283

(1991)), a lipofection method (Proc. Natl. Acad. Sci. USA, 84, 7413 (1987)), and the like.

After introduction of the expression vector, the expression level and antigen binding activity of the humanized antibody in the culture supernatant can be measured by enzyme immunoassay (hereinafter referred to as "ELISA"; *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Chapter 14 (1988), hereinafter referred to as "*Antibodies: A Laboratory Manual*", *Monoclonal Antibodies: Principles and Practice*, Academic Press Limited (1996), hereinafter referred to as "*Monoclonal Antibodies*" (1996)) and the like.

(9) Stable expression of humanized antibody

A transformant which produces a humanized antibody stably can be obtained by introducing into an appropriate host cell the humanized antibody expression vector described in the items 1(4) and 1(7).

Examples of the method for introducing the expression vector into a host cell include electroporation (Japanese Published Unexamined Patent Application No. 257891/90, *Cytotechnology*, 3, 133 (1990)) and the like.

Any cell can be used as the host cell into which the humanized antibody expression vector is to be introduced, so long as it can express a humanized antibody. Examples include mouse SP2/0-Ag14 cell (ATCC CRL1581), mouse P3X63-Ag8.653 cell (ATCC CRL1580), CHO cell in which a dihydrofolate reductase gene (hereinafter referred to as "DHFR gene") is defective (Proc. Natl. Acad. Sci. U.S.A., 77: 4216 (1980)), rat YB2/3HL.P2.G11.16Ag.20 cell (ATCC CRL1662, hereinafter referred to as "YB2/0 cell"), and the like. The YB2/0 cell is preferable, since ADCC activity of the humanized antibody is enhanced when expressed therein.

After introduction of the expression vector, transformants which express a humanized antibody stably are

selected in accordance with the method disclosed in Japanese Published Unexamined Patent Application No. 257891/90, by culturing in a medium for animal cell culture containing an agent such as G418 sulfate (hereinafter referred to as "G418", manufactured by Sigma) or the like. Examples of the medium for animal cell culture include PRMI1640 medium (manufactured by Nissui Pharmaceutical), GIT medium (manufactured by Nissui Pharmaceutical), EX-CELL302 medium (manufactured by JRH), IMDM medium (manufactured by GIBCO BRL), Hybridoma-SFM medium (manufactured by GIBCO BRL), media obtained by adding various additives such as fetal bovine serum (hereinafter referred to as "FBS") to these media, and the like. The humanized antibody can be produced and accumulated in a culture medium by culturing the selected transformants in a medium. The expression level and antigen binding activity of the humanized antibody in the culture supernatant can be measured by ELISA or the like. Also, in the transformant, the expression level of the humanized antibody can be increased by using dhfr amplification system or the like according to the method disclosed in Japanese Published Unexamined Patent Application No. 257891/90.

The humanized antibody can be purified from the culture supernatant of the transformant by using a protein A column (*Antibodies, A Laboratory Manual, Monoclonal Antibodies* (1996)). Any other conventional methods for protein purification can be used. For example, the humanized antibody can be purified by a combination of gel filtration, ion-exchange chromatography, ultrafiltration and the like. The molecular weight of the H chain or the L chain of the purified humanized antibody or the antibody molecule as a whole is determined by polyacrylamide gel electrophoresis (hereinafter referred to as "SDS-PAGE") (Nature, 227: 680 (1970)), Western blotting (*Antibodies, A*

Laboratory Manual, Monoclonal Antibodies (1996)), and the like.

(10) Evaluation of activity of humanized antibody

5 The binding activity to an antigen and the binding activity to cultured cancer cell lines of the purified humanized antibody can be measured by ELISA, an immunofluorescent method (*Cancer Immunol. Immunother.*, 36, 373 (1993)) and the like means. The cytotoxic activity
10 against an antigen positive culture cell line can be evaluated by measuring the CDC activity, the ADCC activity or the like (*Cancer Immunol. Immunother.*, 36, 373 (1993)).

(11) Application method of humanized antibody

15 It is considered to be useful in the diagnosis and treatment of human cancers and the like, such as lung cancer, melanoma, glioma, neuroblastoma, etc., since the humanized antibody specifically binds to GD3 which expresses in cultured cancer cell lines derived from human
20 and shows cytotoxic activities such as CDC activity, ADCC activity and the like. In addition, since the majority of the antibody are derived from the amino acid sequence of a human antibody in comparison with antibodies of non-human animals, it is expected that it will show strong anti-tumor
25 effects in the human body without showing immunogenicity, and the therapeutic effects will be maintained for a prolonged period of time.

 The humanized antibody of the present invention can be administered alone, but it is generally preferred to
30 provide it in the form of a pharmaceutical formulation produced by mixing it with at least one pharmaceutically acceptable carrier in accordance with a method well known in the technical field of pharmaceuticals.

 It is preferred to select a route of administration
35 which is the most effective in carrying out the intended

treatment such as oral administration or parenteral administration, e.g., intraoral administration, tracheal administration, rectal administration, subcutaneous injection, intramuscular injection, intravenous injection, and the like. Intravenous injection is preferred in an antibody formulation.

The dosage form includes sprays, capsules, tablets, granules, syrups, emulsions, suppositories, injections, ointments, tapes, and the like.

Examples of formulation suitable for oral administration include emulsions, syrups, capsules, tablets, powders, granules, and the like.

Liquid preparations such as emulsions and syrups, can be produced using additives such as water, saccharides, e.g., sucrose, sorbitol, fructose, etc., glycols, e.g., polyethylene glycol, propylene glycol, etc., oils, e.g., sesame oil, olive oil, soybean oil, etc., antiseptics, e.g., p-hydroxybenzoate, etc., flavors, e.g., strawberry flavor, peppermint, etc., and the like.

Capsules, tablets, powders, granules and the like can be produced using additives such as fillers, e.g., lactose, glucose, sucrose, mannitol, etc., disintegrating agents, e.g., starch, sodium alginate, etc., lubricants, e.g., magnesium stearate, etc., binders, e.g., polyvinyl alcohol, hydroxypropylcellulose, gelatin, etc., surfactants, e.g., fatty acid ester, etc., plasticizers, e.g., glycerine, etc., and the like.

Examples of formulations suitable for parenteral administration include injections, suppositories, sprays, and the like.

Injections can be prepared using a carrier such as a salt solution, glucose solution or a mixture thereof, or the like.

Suppositories can be prepared using a carrier such as cacao butter, hydrogenated fat, a carboxylic acid, or the like.

Also, sprays can be prepared from the antibody or peptide itself or using a carrier or the like which does not stimulate oral and airway mucous membranes of patients and can facilitate absorption of the antibody or peptide by dispersing it as minute particles.

Examples of the carrier include lactose, glycerine, and the like. Depending on the properties of the antibody or peptide and the carrier to be used, aerosols, dry powders and the like can be produced. The additives exemplified in the oral preparations can also be added to the parenteral preparations.

The dose and frequency of administration vary depending on intended therapeutic effect, administration method, treating period, age, body weight and the like, but the dose is generally from 10 $\mu\text{g/kg}$ to 8 mg/kg per day per adult.

2. Production of fusion protein of antibody or antibody fragment with cytokine

(1) Construction of gene encoding fusion protein of antibody or antibody fragment with cytokine

A gene encoding a fusion protein of an antibody or the antibody fragment with cytokine can be constructed by linking a gene encoding cytokine to the 5'-terminal or 3'-terminal of a gene encoding the H chain or L chain of an antibody or the antibody fragment via an appropriate synthetic DNA. Also, a gene encoding a fusion protein of an antibody or the antibody fragment with cytokine can also be constructed by introducing an appropriate restriction enzyme recognition sequence into the 5'-terminal of a primer for amplification when a gene encoding cytokine is amplified by PCR, and linking the resulting gene to the 5'-

terminal or 3'-terminal of a gene encoding the H chain or L chain of an antibody or the antibody fragment. Any chromosomal DNA or cDNA can be used as the gene encoding cytokine. The nucleotide sequence of the thus constructed gene encoding a fusion protein of an antibody or antibody fragment with cytokine is determined by the method described in the item 1(2) to confirm that it is the sequence of interest.

10 (2) Construction of expression vector of fusion protein of antibody or antibody fragment with cytokine

An expression vector of a fusion protein of an antibody or the antibody fragment with cytokine can be constructed by substituting a part or whole of a gene encoding the H chain or L chain of an antibody or the antibody fragment on the humanized antibody expression vector described in the items 1(4) and 1(7) by the gene encoding a fusion protein of an antibody or the antibody fragment with cytokine described in the item 2(1). For example, when a fusion protein in which cytokine is conjugated with the C-terminal of the H chain of an antibody is prepared, the expression vector can be constructed by linking a gene encoding cytokine to the 3'-terminal of a gene encoding the CH of an antibody or the antibody fragment to thereby construct a gene encoding a fusion protein of the CH of an antibody or the antibody fragment with cytokine according to the item 2(1), and substituting, with the gene, a gene encoding the CH of an antibody or the antibody fragment on the humanized antibody expression vector described in the items 1(4) and 1(7).

(3) Stable expression of fusion protein of antibody or antibody fragment with cytokine

Stable expression of a fusion protein of an antibody or the antibody fragment with cytokine in

accordance with the method described in the item 1(9) using the expression vector of a fusion protein of an antibody or the antibody fragment with cytokine described in the item 2(2) is carried out so that a transformant capable of stably expressing the fusion protein of an antibody or the antibody fragment with cytokine can be obtained, the fusion protein of an antibody or antibody fragment with cytokine is purified from its culture supernatant, and its molecular weight and the like can be analyzed.

(4) Evaluation of *in vitro* activity of fusion protein of antibody or antibody fragment with cytokine

Among activities of the thus purified fusion protein of an antibody or the antibody fragment with cytokine, activities of the antibody moiety, namely binding activity with an antigen and its binding activity to cultured cancer cell lines can be measured by ELISA, an immunofluorescent method and the like. Also, its cytotoxic activities against antigen-positive cultured cancer cell lines can be evaluated by measuring its CDC activity, ADCC activity and the like. On the other hand, activity of the cytokine moiety can be evaluated, e.g., using growth-supporting activity of a cultured cell line which shows concentration-dependent growth against cytokine, as an index (Proc. Natl. Acad. Sci. U.S.A., 91, 9626 (1994)).

(5) Evaluation of *in vivo* activity of fusion protein of an antibody or antibody fragment with cytokine

Antitumor effect of the fusion protein of an antibody or the antibody fragment with cytokine can be evaluated by administering to a syngenic graft mouse model in which an antigen-expressing cultured mouse cancer cell line is transplanted into a wild type mouse having normal immune system. Also, its antitumor effect in the living body can be evaluated by comparing respective cases of the

administration of the antibody alone, cytokine alone and simultaneous administration of the antibody and cytokine to the syngenic graft mouse model (*Proc. Natl. Acad. Sci. U.S.A.*, 93, 7826 (1996)). To date, there are no reports on
5 GD2- or GD3-positive mouse culture cell lines, which are specifically expressed in human melanoma and neuroblastoma. However, a GD2- or GD3-positive transformant can be produced by introducing a ganglioside synthase gene of interest to be specifically expressed into a mouse culture
10 cell line which expresses GM2 or GM3 as a biosynthesis precursor of GD2 or GD3 (*Proc. Natl. Acad. Sci. U.S.A.*, 91, 10455 (1994)). A syngenic graft model can be produced by transplanting this transformant into a wild type mouse having a normal immune system.

15 The thus prepared mouse model can be used in the in vivo evaluation of a fusion protein of an antibody or the antibody fragment with cytokine against a ganglioside of interest.

20 (6) Application method of fusion proteins of humanized antibody or antibody fragment with cytokine

Since the fusion protein of an antibody or the antibody fragment with cytokine of the present invention specifically binds to GD3 expressing in cultured cancer
25 cell lines of human origin and shows cytotoxic activities such as CDC activity, ADCC activity and the like, it is considered to be useful in the diagnosis and treatment of human cancers and the like such as lung cancer, melanoma, glioma, neuroblastoma, etc. Since the majority of the
30 humanized antibody or antibody fragment of the present invention are derived from the amino acid sequence of a human antibody in comparison with antibodies of non-human animals, it is expected that it will show strong anti-tumor effects in the human body without showing immunogenicity,
35 and the therapeutic effects will be maintained for a

prolonged period of time, and since it can activate immunocompetent cells in the periphery of a cancer by the activity of the conjugated cytokine moiety, stronger antitumor effects are expected than the administration of the antibody alone, cytokine alone or simultaneous administration of the antibody and cytokine, and reduction of side effects is expected in comparison with the systemic administration of cytokine.

The fusion protein of a humanized antibody or the antibody fragment with cytokine of the present invention can be administered alone, but it is generally preferred to provide it in the form of a pharmaceutical formulation produced by mixing it with at least one pharmaceutically acceptable carrier in accordance with a method well known in the technical field of pharmaceuticals.

It is preferred to select a route of administration which is the most effective in carrying out the intended treatment such as oral administration or parenteral administration, e.g., intraoral administration, tracheal administration, rectal administration, subcutaneous injection, intramuscular injection, intravenous injection, and the like. Intravenous injection is preferred in a protein formulation.

The dosage form includes sprays, capsules, tablets, granules, syrups, emulsions, suppositories, injections, ointments, tapes, and the like.

Examples of formulation suitable for oral administration include emulsions, syrups, capsules, tablets, powders, granules, and the like.

Liquid preparations such as emulsions and syrups, can be produced using additives such as water, saccharides, e.g., sucrose, sorbitol, fructose, etc., glycols, e.g., polyethylene glycol, propylene glycol, etc., oils, e.g., sesame oil, olive oil, soybean oil, etc., antiseptics, e.g.,

p-hydroxybenzoate, etc., flavors, e.g., strawberry flavor, peppermint, etc., and the like.

Capsules, tablets, powders, granules and the like can be produced using additives such as fillers, e.g.,
5 lactose, glucose, sucrose, mannitol, etc., disintegrating agents, e.g., starch, sodium alginate, etc., lubricants, e.g., magnesium stearate, etc., binders, e.g., polyvinyl alcohol, hydroxypropylcellulose, gelatin, etc., surfactants, e.g., fatty acid ester, etc., plasticizers, e.g., glycerine,
10 etc., and the like.

Examples of formulations suitable for parenteral administration include injections, suppositories, sprays, and the like.

Injections can be prepared using a carrier such as
15 a salt solution, glucose solution or a mixture thereof, or the like.

Suppositories can be prepared using a carrier such as cacao butter, hydrogenated fat, a carboxylic acid, or the like.

20 Also, sprays can be prepared from the antibody or peptide itself or using a carrier or the like which does not stimulate oral and airway mucous membranes of patients and can facilitate absorption of the antibody or peptide by dispersing it as minute particles.

25 Examples of the carrier include lactose, glycerine, and the like. Depending on the properties of the antibody or peptide and the carrier to be used, aerosols, dry powders and the like can be produced. The additives exemplified in the oral preparations can also be added to
30 the parenteral preparations.

The dose and frequency of administration vary depending on intended therapeutic effect, administration method, treating period, age, body weight and the like, but the dose is generally from 10 $\mu\text{g/kg}$ to 8 mg/kg per day per
35 adult.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a drawing showing construction steps of plasmid phKM641H.

5 Fig. 2 is a drawing showing construction steps of plasmid phKM641L.

Fig. 3 is a drawing showing construction steps of plasmid pT796H641L.

10 Fig. 4 is a drawing showing construction steps of plasmid pBS641R.

Fig. 5 is a drawing showing construction steps of plasmid pT641.

Fig. 6 is a drawing showing construction steps of plasmid pT641HCDR.

15 Fig. 7 is a drawing showing construction steps of plasmids pT641LCDR and pT641HLCDR.

Fig. 8 is a drawing showing activity evaluation of an anti-GD3 chimeric antibody and an anti-GD3 CDR-grafted antibody by their transient expression using plasmids pT641, 20 pT641HCDR, pT641LCDR and pT641HLCDR. The ordinate and the abscissa represent the expression vector name and the relative activity (%) when the activity of anti-GD3 chimeric antibody is defined as 100, respectively.

25 Fig. 9 is a drawing showing construction steps of plasmid phKM641Lm-69.

Fig. 10 is a drawing showing construction steps of plasmids pT641HLCDRNL, pT641HLCDRm-1, pT641HLCDRm-4, 30 pT641HLCDRm-6, pT641HLCDRm-7, pT641HLCDRm-8, pT641HLCDRm-9 and pT641HLCDRm-69.

30 Fig. 11 is a drawing showing activity evaluation of anti-GD3 chimeric antibody and anti-GD3 CDR-grafted antibody by their transient expression using plasmids pT641, pT641HLCDR, pT641HLCDRNL, pT641HLCDRm-1, pT641HLCDRm-4, 35 pT641HLCDRm-6, pT641HLCDRm-7, pT641HLCDRm-8, pT641HLCDRm-9 and pT641HLCDRm-69. The ordinate and the

abscissa represent the expression vector name and the relative activity (%) when the activity of anti-GD3 chimeric antibody is defined as 100, respectively.

Fig. 12 is a drawing showing construction steps of plasmids pKANTEK641HLCDR, pKANTEK641HLCDRIm-9 and pKANTEK641HLCDRIm-69.

Fig. 13 is a drawing showing SDS-PAGE (using a 4 to 15% gradient gel) electrophoresis patterns of purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871. The left side shows results of electrophoresis carried out under non-reducing conditions, and the right side shows those under reducing conditions. Lanes 1, 2, 3, 4, 5, 6, 7 and 8 show electrophoresis patterns of high molecular weight markers, KM8869, KM8870, KM8871, low molecular weight markers, KM8869, KM8870 and KM8871, respectively.

Fig. 14 is a drawing showing binding activity of purified anti-GD3 chimeric antibody KM871 and purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 to GD3 measured by changing the antibody concentration. The ordinate and the abscissa are the binding activity to GD3 and the antibody concentration, respectively. "○", "●", "□" and "■" show the activities of KM871, KM8869, KM8870 and KM8871, respectively.

Fig. 15 is a drawing showing binding activity of purified anti-GD3 chimeric antibody KM871 and purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 to GD3 measured by changing an amount of GD3 to be adsorbed to a plate. The ordinate and the abscissa are the binding activity to GD3 and the amount of GD3 adsorbed to the plate, respectively. "○", "●", "□" and "■" show the activities of KM871, KM8869, KM8870 and KM8871, respectively.

Fig. 16 is a drawing showing reactivity of purified anti-GD3 chimeric antibody KM871 and purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 with various

gangliosides. The ordinate and the abscissa are the kind of ganglioside and the binding activity, respectively. AcGM2, GcGM2, AcGM3 and GcGM3 indicate N-acetylGM2, N-glycolylGM2, N-acetylGM3 and N-glycolylGM3. "□", "□", "□" and "■" show the reactivities of KM871, KM8869, KM8870 and KM8871, respectively.

Fig. 17 is a drawing showing reactivity of purified anti-GD3 chimeric antibody KM871 and purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 with human melanoma cell lines G-361 and SK-MEL-28. The ordinate and the abscissa are the number of cells and the fluorescence intensity, respectively. The graphs show reactivities of control, KM871, KM8869, KM8870 and KM8871, respectively, from the bottom.

Fig. 18 is a drawing showing CDC activity of purified anti-GD3 chimeric antibody KM871 and purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 against human melanoma cell lines G-361 and SK-MEL-28. The ordinate and the abscissa are the cytotoxic activity and the antibody concentration, respectively. "□", "□", "□" and "■" show the reactivities of KM871, KM8869, KM8870 and KM8871, respectively.

Fig. 19 is a drawing showing ADCC activity of purified anti-GD3 chimeric antibody KM871 and purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 against human melanoma cell lines G-361 and SK-MEL-28. The ordinate and the abscissa are the cytotoxic activity and the antibody concentration, respectively. "□", "□", "□" and "■" show the reactivities of KM871, KM8869, KM8870 and KM8871, respectively.

Fig. 20 is a drawing showing construction steps of plasmid pBSA-B.

Fig. 21 is a drawing showing construction steps of plasmid pBSAhCyl-IL-2.

Fig. 22 is a drawing showing construction steps of plasmid pBShCyl-IL-2.

Fig. 23 is a drawing showing construction steps of plasmid pKANTEX8871-hIL-2.

5 Fig. 24 is a drawing showing SDS-PAGE (using a 4 to 15% gradient gel) electrophoresis patterns of purified anti-GD3 CDR-grafted antibody KM8871 and purified fusion protein KM8871-hIL-2. The left side and the right side are results of electrophoresis carried out under non-reducing conditions, and those under reducing conditions, respectively. Lanes 1, 2, 3, 4, 5 and 6 show electrophoresis patterns of low molecular weight markers, KM8871, KM8871-hIL-2, low molecular weight markers, KM8871 and KM8871-hIL-2, respectively.

10 Fig. 25 is a drawing showing binding activity of purified anti-GD3 CDR-grafted antibody KM8871 and purified fusion protein KM8871-hIL-2 to GD3 measured by changing the antibody concentration. The ordinate and the abscissa are the binding activity to GD3 and the antibody concentration. "○" and "●" show the activities of KM8871 and of KM8871-hIL-2, respectively.

15 Fig. 26 is a drawing showing reactivity of purified anti-GD3 CDR-grafted antibody KM8871 and purified fusion protein KM8871-hIL-2 with various gangliosides. The ordinate and the abscissa are the kind of ganglioside and the binding activity. AcGM2, GcGM2, AcGM3 and GcGM3 indicate N-acetylGM2, N-glycolylGM2, N-acetylGM3 and N-glycolylGM3, respectively. "□" and "■" show the reactivities of KM8871 and KM8871-hIL-2, respectively.

25 Fig. 27 is a drawing showing reactivity of purified anti-GD3 CDR-grafted antibody KM8871 and purified fusion protein KM8871-hIL-2 with human melanoma cell line G-361. The ordinate and the abscissa are the number of cells and the fluorescence intensity, respectively. The drawings

show the reactivities of KM8871 and KM8871-hIL-2, respectively, from the upper column.

Fig. 28 is a drawing showing growth-supporting activity of hIL-2 and purified fusion protein KM8871-hIL-2 against hIL-1-dependent cell CTLL-2 measured by changing concentration of each protein. The ordinate and the abscissa are the growth-supporting activity and the protein concentration, respectively. "○" and "●" indicate the activities of hIL-2 and KM8871-hIL-2, respectively.

Fig. 29 is a drawing showing results of the measurement of the activation and accompanying cytotoxic activity of human effector cells by purified anti-GD3 CDR-grafted antibody KM8871 and purified fusion protein KM8871-hIL-2. The ordinate and the abscissa are the cytotoxic activity and the used protein, respectively. "□" and "■" indicate the activities of KM8871 and KM8871-hIL-2, respectively.

Fig. 30 is a drawing showing construction steps of plasmid pKANTEX871-hIL-2.

Fig. 31 is a drawing showing SDS-PAGE (using a 4 to 15% gradient gel) electrophoresis patterns of anti-GD3 chimeric antibody KM871 and purified fusion protein KM871-hIL-2. The left side and the right side are results of electrophoresis carried out under reducing conditions and under non-reducing conditions, respectively. Lanes 1, 2, 3, 4, 5 and 6 show electrophoresis patterns of low molecular weight markers, KM871, KM871-hIL-2, low molecular weight markers, KM871 and KM871-hIL-2, respectively.

Fig. 32 is a drawing showing binding activity of anti-GD3 chimeric antibody KM8871 and purified fusion protein KM871-hIL-2 to GD3 measured by changing the antibody concentration. The ordinate and the abscissa are the binding activity to GD3 and the antibody concentration, respectively. "●" and "□" show the activities of KM871 and KM871-hIL-2, respectively.

Fig. 33 is a drawing showing reactivity of anti-GD3 chimeric antibody KM871 and purified fusion protein KM871-hIL-2 with various gangliosides. The ordinate and the abscissa are the kind of ganglioside and the binding activity, respectively. AcGM2, GcGM2, AcGM3 and GcGM3 indicate N-acetylGM2, N-glycolylGM2, N-acetylGM3 and N-glycolylGM3, respectively. "■" and "□" show the reactivities of KM871 and KM871-hIL-2, respectively.

Fig. 34 is a drawing showing reactivity of anti-GD3 chimeric antibody KM871 and purified fusion protein KM871-hIL-2 with human melanoma cell lines G-361 and SK-MEL-28. The ordinate and the abscissa are the number of cells and the fluorescence intensity, respectively. The graphs show the reactivities of KM871 to G361 cell, KM871-hIL-2 to G361 cell, KM871 to SK-MEL-28 and KM871-hIL-2 to SK-MEL-28, respectively, from the upper column.

Fig. 35 is a drawing showing growth-supporting activity of hIL-2 and purified fusion protein KM871-hIL-2 against hIL-2-dependent cell CTLL-2 measured by changing concentration of each protein. The ordinate and the abscissa are the growth-supporting activity and the protein concentration, respectively. "●" and "○" indicate the activities of hIL-2 and KM871-hIL-2, respectively.

Fig. 36 is a drawing showing results of the measurement of the activation and accompanying cytotoxic activity of human effector cells by anti-GD3 chimeric antibody KM871 and purified fusion protein KM871-hIL-2. The ordinate and the abscissa are the cytotoxic activity and the used protein, respectively. "■" and "□" indicate activities of KM871 and KM871-hIL-2, respectively.

Fig. 37 is a drawing showing construction steps of plasmid pKANTEXGD3.

Fig. 38 is a drawing showing reactivity of anti-GD3 chimeric antibody KM871 with GD3 synthase gene-introduced B16 cell lines B16-29-10 and B16-8-3 and the parent line

B16 F10 cell. The ordinate and the abscissa are the number of cells and the fluorescence intensity, respectively. The drawings show the reactivities of B16-F10 cell, B16-29-10 and B16 8-3, respectively, from the upper column.

5 Fig. 39 is a drawing showing results of the measurement of anti-metastatic effects of anti-GD3 chimeric antibody KM871 and KM871-hIL-2 on BL/6 mice into which a GD3-positive B16 cell line B16-29-10 is transplanted. The ordinate and the bar of the graph show the number of
10 metastatic foci on the lung surface and the average value of each group, respectively. No administration means a control group in which the antibody is not administered. " < ", " ○ " and " ● " indicate the numbers of lung-metastasized foci in mice of the control group, the KM871-
15 hIL-2-administered group and the KM871-administered group, respectively.

 Fig. 40 is a drawing showing a result of the measurement of anti-metastatic effects of anti-GD3 chimeric antibody KM871 and KM871-hIL-2 on BL/6 mice into which a
20 GD3-positive B16 cell line B16-8-3 is transplanted. The ordinate and the bar of the graph show the number of metastatic foci on the lung surface and the average value of each group, respectively. No administration means a control group in which the antibody is not administered.
25 " ◇ ", " ○ " and " ● " indicate the numbers of lung-metastasized foci in mice of the control group, the KM871-hIL-2-administered group and the KM871-administered group, respectively.

 Fig. 41 is a drawing showing results of the
30 measurement of anti-metastatic effects by combined administration of anti-GD3 chimeric antibody KM871 and hIL-2 and administration of KM871-hIL-2 on BL/6 mice into which a GD3-positive B16 cell line B16-29-10 is transplanted. The ordinate and the bar of the graph show the number of
35 metastatic foci on the lung surface and the average value

of each group. No administration means a control group in which the antibody is not administered. "◇", "○" and "●" indicate the number of lung-metastasized foci in mice of the control group, the KM871-hIL-2-administered group and the KM871/hIL-2 combined administered group, respectively.

Fig. 42 is a drawing showing results of the measurement of growth inhibition effects of KM871-hIL-2 on a solid tumor early stage model of BL/6 mice into which a GD3-positive B16 cell line B16-29-10 is subcutaneously transplanted. The ordinate and the abscissa of the graph show the ratio of the average value of tumor volumes in each group to the average value of tumor volumes in the control group and days after commencement of the agent administration, respectively. Dotted line, "□", "△" and "○" indicate control, the KM871-administered group, the hIL-2-administered group and the KM871-hIL-2-administered group, respectively.

Fig. 43 is a drawing showing results of the measurement of growth inhibition effects of KM871-hIL-2 on a solid tumor advanced stage model of BL/6 mice into which a GD3-positive B16 cell line B16-29-10 is subcutaneously transplanted. The ordinate of the graph shows the ratio of average value of V/V_0 values in each group to average value of V/V_0 values in the control group. V/V_0 indicates the ratio of the tumor volume of each mouse on each measured day to the tumor volume on the day of commencement of the administration. The abscissa shows days after commencement of the agent administration. Dotted line, "△", "□", "■", "○" and "●" indicate control, the hIL-2-administered group, the KM871-administered group at 50 $\mu\text{g/day}$, the KM871-administered group at 200 $\mu\text{g/day}$, the KM871-hIL-2-administered group at 24 $\mu\text{g/day}$ and KM871-hIL-2-administered group at 60 $\mu\text{g/day}$.

Fig. 44 is a drawing showing results of the measurement of growth inhibition effects by combined

administration of anti-GD3 chimeric antibody KM871 and KM871-hIL-2 on a solid tumor advanced stage model of BL/6 mice into which a GD3-positive B16 cell line B16-29-10 is subcutaneously transplanted. The ordinate of the graph shows the ratio of the average value of V/V0 values in each group to the average value of V/V0 values in the control group. V/V0 indicates the ratio of tumor volume of each mouse on each measured day to tumor volume on the day of commencement of the administration. The abscissa shows days after commencement of the agent administration. Dotted line, " Δ ", " \blacksquare ", " \bigcirc " and " \bullet " indicate control, the hIL-2-administered group, the KM871-administered group, the combined administration group of KM871 and KM871-hIL-2 at 15 μ g/day and the combined administration group of KM871 and KM871-hIL-2 at 60 μ g/day.

Fig. 45 is a drawing showing results of measurement of the activation and accompanying enhance in cytotoxic activity of BL/6 mouse spleen cells by hIL-2 or KM871-hIL-2. The ordinate and the abscissa are the cytotoxic activity and the concentration of KM871 used in the cytotoxic activity measurement, respectively. Black bar, white bar, gray bar and shaded bar indicate the activity when spleen cells of non-administered control mice are used, the activity when spleen cells of mice to which hIL-2 is administered five times are used, the activity when spleen cells of mice to which KM871-hIL-2 is administered once are used and the activity when spleen cells of mice to which KM871-hIL-2 is administered five times are used, respectively.

Fig. 46 is a drawing showing the ratio of the number of cells of each cell group in spleen cells of BL/6 mouse to which hIL-2 or KM871-hIL-2 is administered, measured by an immunofluorescent method. The ordinate and the abscissa are the ratio in spleen cells of each cell group and each cell group, respectively. White bar, gray

bar, black bar and shaded bar indicate the ratio in spleen cells of non-administered control mice, the ratio in spleen cells of mice to which hIL-2 is administered five times, the ratio in spleen cells of mice to which KM871-hIL-2 is administered five times and the ratio in spleen cells of mice to which KM871-hIL-2 is administered twice, respectively.

Fig. 47 is a drawing showing a result of measurement of cytotoxic activity by spleen cells when KM871-hIL-2 is administered to BL/6 mice from which each of NK cell, CD4-positive T cell and CD8-positive T cells are depleted. The ordinate shows cytotoxic activity. White bar, black bar, gray bar and shaded bar indicate the cytotoxic activity of mouse spleen cells which is not subjected to the removing treatment, the cytotoxic activity of NK cell-depleted mouse spleen cells, the cytotoxic activity of CD4-positive T cell-depleted mouse spleen cells and the cytotoxic activity of CD8-positive T cell-depleted mouse spleen cells, respectively.

Fig. 48 is a drawing showing results of the measurement of anti-metastatic effects of KM871-hIL-2 on BL/6 mice in which a GD3-positive B16 cell line B16-29-10 is transplanted into mice from which each of NK cell, CD4-positive T cell and CD8-positive T cell are depleted. The ordinate shows the number of metastatic foci. "○", "●", horizontal bar and the value in the graph indicate the number of metastatic foci of mouse lung, the number of metastatic foci of KM871-hIL-2-administered mouse lung, the average value of each group and the metastasis inhibition ratio by KM871-hIL-2 administration.

BEST MODE FOR CARRYING OUT THE INVENTION

Examples of the present invention are shown below, though the scope of the present invention is not restricted thereby.

Example 1

Production of anti-GD3 CDR-grafted antibody:

1. Analysis of amino acid sequence of V region of anti-GD3
5 mouse antibody KM641

A full amino acid sequence of VH of anti-GD3 mouse
antibody KM641 disclosed in Japanese Published Unexamined
Patent Application No. 304989/93 is represented by SEQ ID
NO:1, and a full amino acid sequence of VL thereof is
10 represented by SEQ ID NO:2. Based on the comparison of
both sequences with analytical results of amino acid
sequences of known antibodies (*Sequences of Proteins of
Immunological Interest*) and N-terminal amino acid sequences
of H chain and L chain of purified anti-GD3 mouse antibody
15 KM641, it was confirmed that positions (-19) to (-1) in the
amino acid sequence of H chain and positions (-20) to (-1)
in the amino acid sequence of L chain are secretory signal
sequences. Full amino acid sequences of the secretory VH
and VL are represented by SEQ ID NO:55 and SEQ ID NO:56,
20 respectively. Also, it was found that CDRs 1, 2 and 3 of
VH have the amino acid sequences represented by SEQ ID
NOS:3, 4 and 5, and CDRs 1, 2 and 3 of VL have those
represented by SEQ ID NOS:6, 7 and 8.

25 2. Measurement of binding activity of antibody to various
ganglioside (ELISA)

The binding activity of antibodies to various
gangliosides was measured as follows.

Each ganglioside (2 nmol) was dissolved in 2 ml of
30 ethanol solution containing 10 µg of
dipalmitoylphosphatidylcholine (manufactured by SIGMA) and
5 µg of cholesterol (manufactured by SIGMA). In each well
of a 96 well plate for ELISA use (manufactured by Greiner),
20 µl of the solution (becomes 20 pmol/well) or 20 µl of
35 the solution diluted with ethanol was respectively

dispensed, followed by air-drying, and then PBS containing 1% BSA (hereinafter referred to as "1% BSA-PBS") was added in 100 µl/well and allowed to react at room temperature for 1 hour to thereby block remaining active groups. After discarding 1% BSA-PBS, culture supernatants of transformants or diluted solutions of humanized antibodies were added in 50 µl/well and allowed to react at room temperature for 1 hour. After the reaction, each well was washed with PBS containing 0.05% Tween 20 (hereinafter referred to as "Tween-PBS") and then a peroxidase-labeled goat anti-human IgG(γ) antibody solution (manufactured by Kirkegaard & Perry Laboratories) diluted 1,000 folds with 1% BSA-PBS was added as a secondary antibody solution in 50 µl/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, an ABTS substrate solution (a solution prepared by dissolving 0.55 g of 2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid) in 1 L of 0.1 M citrate buffer (pH 4.2) and adding 1 µl/ml hydrogen peroxide just before the use) was added in 50 µl/well for color development and the absorbance at 415 nm (hereinafter referred to as "OD415") was measured.

3. Construction of cDNAs encoding VH and VL of anti-GD3 CDR-grafted antibody

(1) Design of amino acid sequences of VH and VL of anti-GD3 CDR-grafted antibody

First, an amino acid sequence of VH of an anti-GD3 CDR-grafted antibody was designed as follows. In order to graft the CDR amino acid sequences of VH of the anti-GD3 mouse antibody KM641 identified in the item 1 of Example 1, an FR amino acid sequence of VH of a human antibody was selected. Kabat et al. have classified the VH of various known human antibodies into three subgroups (HSG I to III) based on the homology of the amino acid sequences and then

reported consensus sequences among respective subgroups
(Sequences of Proteins of Immunological Interest). Since
these consensus sequences have a possibility that the
immunogenicity is reduced in human, it was decided to
5 design a VH amino acid sequence of an anti-GD3 CDR-grafted
antibody based on these consensus sequences. In order to
produce an anti-GD3 CDR-grafted antibody having higher
activity in designing it, it was decided to select an FR
amino acid sequence having the highest homology with the FR
10 amino acid sequence of VH of KM641, among FR amino acid
sequences of consensus sequences of the three subgroups of
VH of human antibodies. Results of the homology search are
shown in Table 1. As shown in Table 1, the FR amino acid
sequence of VH of KM641 showed the most high homology with
15 the subgroup III.

Table 1
Homology between FR amino acid sequence of
consensus sequence of each subgroup of
20 human antibody H chain V region and
FR amino acid sequence of H chain V region of KM641

HSG I	HSG II	HSG III
62.1%	56.3%	78.2%

Based on the above results, an amino acid sequence
25 HV.0 of VH of the anti-GD3 CDR-grafted antibody was
designed by grafting the CDR amino acid sequence of VH of
the anti-GD3 mouse antibody KM641 to an appropriate
position of the amino acid sequence of FR of the consensus
sequence of subgroup III of VH of the human antibody.

30 Next, an amino acid sequence of VL of an anti-GD3
CDR-grafted antibody was designed as follows. In order to
graft the CDR amino acid sequence of VL of the anti-GD3

mouse antibody KM641 identified in the item 1 of Example 1,
an FR amino acid sequence of VL of a human antibody was
selected. Kabat et al. have classified the VL of various
known human antibodies into four subgroups (HSG I to IV)
5 based on the homology of the amino acid sequences and then
reported consensus sequences among respective subgroups
(*Sequences of Proteins of Immunological Interest*).
Accordingly, similar to the H chain, an FR amino acid
sequence having most high homology with the FR amino acid
10 sequence of VL of KM641 was selected from FR amino acid
sequences of consensus sequences of the four subgroups of
VL of human antibodies. Results of the homology search are
shown in Table 2. As shown in Table 2, the FR amino acid
sequence of VL of KM641 showed the most high homology with
15 the subgroup I.

Table 2
Homology between FR amino acid sequence of
consensus sequence of each subgroup of
human antibody L chain V region and
20 FR amino acid sequence of L chain V region of KM641

HSG I	HSG II	HSG III	HSG IV
76.2%	60.0%	62.5%	67.5%

Based on the above results, an amino acid sequence
25 LV.0 of VL of the anti-GD3 CDR-grafted antibody was
designed by grafting the CDR amino acid sequence of VL of
the anti-GD3 mouse antibody KM641 to an appropriate
position of the amino acid sequence of FR of the consensus
sequence of subgroup I of VL of the human antibody.

30 The thus designed amino acid sequence HV.0 of VH
and amino acid sequence LV.0 of VL of the anti-GD3 CDR-
grafted antibody are sequences in which only the CDR amino

acid sequence of the anti-GD3 mouse antibody is grafted to the FR amino acid sequence of the selected human antibody. In general, in the human CDR-grafted antibodies, the antigen binding activity is reduced in many cases by grafting of a mouse antibody CDR amino acid sequence alone. Thus, in order to avoid the reduction of the antigen binding activity of antibodies, among FR amino acid residues different between a human antibody and a mouse antibody, amino acid residues considered to be exerting influences on the antigen binding activity are grafted together with a CDR amino acid sequence. Accordingly, identification of FR amino acid residues considered to have influences on the activity was carried out.

First, a three-dimensional structure of an antibody V region comprised of the amino acid sequence HV.0 of VH and amino acid sequence LV.0 of VL of anti-GD3 CDR-grafted antibody designed in the above (hereinafter referred to as "HV0LV0") was constructed using computer-modeling. The production of three-dimensional structure coordinates was carried out using a software AbM (manufactured by Oxford Molecular) and the display of three-dimensional structure was carried out using a software Pro-Explore (manufactured by Oxford Molecular) in accordance with the respective manufacture's instructions. Also, a computer model of the three-dimensional structure of the V region of anti-GD3 mouse antibody KM641 was constructed in the same manner. In addition, a three-dimensional structure model of a modified HV0LV0 having an amino acid sequence in which at least one amino acid residue different from anti-GD3 mouse antibody KM641 in the FR amino acid sequences of VH and VL of HV0LV0 was substituted by the amino acid residues of positions corresponding to the anti-GD3 mouse antibody KM641 in order was also constructed. Three-dimensional structures of V regions of the anti-GD3 mouse antibody KM641, HV0LV0 and modified product were compared. As a

result, amino acid residues considered to have influences on the antigen binding activity by changing three-dimensional structure of the antigen-binding region were selected from the FR amino acid residues of HV0LV0. As a result of substituting the thus selected FR amino acid residues of HV0LV0 by the amino acid residues found in the mouse antibody KM641, an amino acid sequence hKM641H of VH of the anti-GD3 CDR-grafted antibody represented by SEQ ID NO:9 and an amino acid sequence hKM641L of VL of the anti-GD3 CDR-grafted antibody represented by SEQ ID NO:10 were designed. In the hKM641H, the 10th position Gly, the 11th position Leu, the 20th position Leu, the 28th position Thr, the 84th position Asn, the 91st position Thr, the 95th position Tyr, the 97th position Ala and the 115th position Val in the FR amino acid sequence of HV.0 were replaced by Asp, Phe, Val, Ala, Arg, Ser, Phe, Thr and Leu, respectively, as amino acid residues of positions corresponding to the VH of anti-GD3 mouse antibody KM641. In the hKM641L, the 49th position Tyr, the 65th position Ser and the 71st position Phe in the FR amino acid sequence of LV.0 were replaced by Tyr, Ser and Phe, respectively, as amino acid residues of positions corresponding to the VL of anti-GD3 mouse antibody KM641.

(2) Construction of cDNA encoding VH of anti-GD3 CDR-grafted antibody

A cDNA encoding the anti-GD3 CDR-grafted antibody VH amino acid sequence hKM641H designed in the item 3(1) of Example 1 was constructed as follows.

First, the designed amino acid sequence was ligated with the secretory signal sequence of H chain of anti-GD3 mouse antibody KM641 represented by SEQ ID NO:1 to produce a full antibody amino acid sequence. Next, the amino acid sequence is converted into genetic codons. When two or more genetic codons were present for one amino acid residue,

corresponding genetic codon was determined by taking the frequency of codon usage found in nucleotide sequences of antibody genes into consideration (*Sequences of Proteins of Immunological Interest*). A nucleotide sequence of cDNA
5 encoding the complete antibody V region amino acid sequence was designed by ligating the thus determined genetic codons, and nucleotide sequences of primer binding site for use in the PCR amplification (including restriction enzyme recognition sequences for cloning into a humanized antibody
10 expression vector) were added to the 5'-terminal and 3'-terminal. The thus designed nucleotide sequence was divided into a total of 6 nucleotide sequences from the 5'-terminal side, each having about 100 bases (adjoining nucleotide sequences are designed such that the termini
15 have an overlapping sequence of about 20 bases), and they were synthesized in alternating order of the sense chain and the antisense chain using an automatic DNA synthesizer (380A, manufactured by Applied Biosystems).

Specifically, 6 synthetic DNA fragments of SEQ ID
20 NO:11 to SEQ ID NO:16 were synthesized. Each DNA was added to 50 μ l of a buffer comprising 10 mM Tris-HCl (pH 8.3), 50 mM potassium chloride, 1.5 mM magnesium chloride, 0.001% gelatin, 200 μ M dNTPs, 0.5 μ M M13 primer RV (manufactured by Takara Shuzo), 0.5 μ M M13 primer M4 (manufactured by
25 Takara Shuzo) and 2 units of TaKaRa Taq DNA polymerase (manufactured by Takara Shuzo) to give a final concentration of 0.1 μ M, and the solution was covered with 50 μ l of mineral oil and set to a DNA thermal cycler (PJ480, manufactured by PERKIN ELMER) to carry out 30 cycles of the
30 reaction, each cycle including 2 minutes at 94°C, 2 minutes at 55°C and 2 minutes at 72°C. The reaction solution was purified using QIAquick PCR Purification Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions and made into 30 μ l of a buffer comprising 10
35 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT,

and 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT, 100 µg/ml BSA and 0.01% Triton X-100, and 10 units of a restriction enzyme *NotI* (manufactured by Takara Shuzo) were further added to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 0.2 µg of an *ApaI-NotI* fragment of about 0.44 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 3 µg of a plasmid pBluescript SK(-) was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol and added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT, 100 µg/ml BSA and 0.01% Triton X-100, and 10 units of a restriction enzyme *NotI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 2 µg of an *ApaI-NotI* fragment of about 2.95 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 0.1 µg of the *ApaI-NotI* fragment of the PCR product of VH of anti-GD3 CDR-grafted antibody and 0.1 µg of the *ApaI-NotI* fragment of the plasmid pBluescript SK(-), both obtained in the above, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus

obtained recombinant plasmid DNA solution, an *Escherichia coli* line HB101 was transformed. Each plasmid DNA was produced from 10 clones of the transformants, allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence, and as a result, the plasmid pHKM641H shown in Fig. 1 having the nucleotide sequence of interest was obtained.

(3) Construction of cDNA encoding VL of anti-GD3 CDR-grafted antibody

A cDNA encoding the anti-GD3 CDR-grafted antibody VL amino acid sequence hKM641L designed in the item 3(1) of Example 1 was constructed as follows using the PCR similar to the VH. In this case, the amino acid sequence of L chain of anti-GD3 mouse antibody KM641 represented by SEQ ID NO:2 was used as the secretory signal sequence.

First, 6 synthetic DNA fragments of SEQ ID NO:17 to SEQ ID NO:22 were synthesized using an automatic DNA synthesizer (380A, manufactured by Applied Biosystems). Each of the thus synthesized DNA fragments was added to 50 μ l of a buffer comprising 10 mM Tris-HCl (pH 8.3), 50 mM potassium chloride, 1.5 mM magnesium chloride, 0.001% gelatin, 200 μ M dNTPs, 0.5 μ M M13 primer RV (manufactured by Takara Shuzo), 0.5 μ M M13 primer M4 (manufactured by Takara Shuzo) and 2 units of TaKaRa Taq DNA polymerase (manufactured by Takara Shuzo) to give a final concentration of 0.1 μ M, and the solution was covered with 50 μ l of mineral oil and set to a DNA thermal cycler (PJ480, manufactured by PERKIN ELMER) to carry out 30 cycles of the reaction, each cycle including 2 minutes at 94°C, 2 minutes at 55°C and 2 minutes at 72°C. The reaction solution was purified using QIAquick PCR Purification Kit (manufactured

by QIAGEN) in accordance with the manufacture's instructions and made into 30 μ l of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 μ g/ml BSA, and 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) and a restriction enzyme *SplI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 0.2 μ g of an *EcoRI-SplI* fragment of about 0.39 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 3 μ g of a plasmid pBSL3 described in Japanese Published Unexamined Patent Application No. 257893/98 was added to 10 μ l of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 μ g/ml BSA, and 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) and a restriction enzyme *SplI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 2 μ g of an *EcoRI-SplI* fragment of about 2.95 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 0.1 μ g of the *EcoRI-SplI* fragment of the PCR product of VL of anti-GD3 CDR-grafted antibody and 0.1 μ g of the *EcoRI-SplI* fragment of the plasmid pBSL3, both obtained in the above, were added to 20 μ l in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained recombinant plasmid DNA solution, an *E. coli* line HB101 was transformed. Each plasmid DNA was prepared from 10 clones of the transformants, allowed to react in accordance with the manufacture's instructions attached to

AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence, and as a result, the plasmid pHKM641L shown in Fig. 2 having the nucleotide sequence of interest was obtained.

4. Activity evaluation of anti-GD3 CDR-grafted antibody by transient expression using animal cell

In order to carry out activity evaluation of anti-GD3 CDR-grafted antibody more quickly, transient expression of anti-GD3 CDR-grafted antibody was carried out as follows using COS-7 cell (ATCC CRL 1651).

(1) Construction of anti-GD3 chimeric antibody KM871 transient expression vector pT641

In order to use as a positive control in the activity evaluation of anti-GD3 CDR-grafted antibody by transient expression, a transient expression vector pT641 for anti-GD3 chimeric antibody KM871 was constructed as follows.

Since the efficiency of transient expression using animal cells generally depends on the copy number of introduced expression vector, it was considered that smaller expression vector has higher expression efficiency. Accordingly, the transient expression vector pT641 for anti-GD3 chimeric antibody KM871 was constructed as follows using the transient expression vector pT796 for anti-ganglioside GM2 chimeric antibody KM966 described in Japanese Published Unexamined Patent Application No. 257893/98 and the plasmid pKM641HF1 having VH of anti-GD3 mouse antibody KM641 and plasmid pKM641LA2 having VL of anti-GD3 mouse antibody KM641 described in Japanese Published Unexamined Patent Application No. 304989/93.

First, 3 µg of the plasmid pKM641LA2 having VL of anti-GD3 mouse antibody KM641 described in Japanese Published Unexamined Patent Application No. 304989/93 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 50 mM sodium chloride, 10 mM magnesium chloride and 1 mM DTT and the solution was further mixed with 10 units of a restriction enzyme *HindIII* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol and added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride and 1 mM DTT, and 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) were further added to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 0.2 µg of an *HindIII-EcoRI* fragment of about 0.35 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 3 µg of the transient expression vector pT796 for anti-ganglioside GM2 chimeric antibody KM966 described in Japanese Published Unexamined Patent Application No. 257893/98 was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 µg/ml BSA, and the solution was further mixed with 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) and a restriction enzyme *SplI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 2 µg of an *EcoRI-SplI* fragment of about 9.20 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, synthetic DNA fragments respectively having the nucleotide sequences represented by SEQ ID NO:23 and SEQ ID NO:24 were synthesized using an automatic DNA

synthesizer (380A, manufactured by Applied Biosystems). A 0.3 µg of each of the thus obtained synthetic DNA fragments was added to 15 µl of sterile water and heated at 65°C for 5 minutes. After allowing the reaction solution to stand at room temperature for 30 minutes, this was mixed with 2 µl of a 10 × buffer solution (500 mM Tris-HCl (pH 7.6), 100 mM magnesium chloride, 50 mM DTT) and 2 µl of 10 mM ATP and further mixed with 10 units of T4 Polynucleotide Kinase (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 30 minutes to thereby phosphorylate the 5'-terminal.

Next, 0.1 µg of the *Hind*III-*Eco*RI fragment derived from the plasmid pKM641LA2, 0.1 µg of the *Eco*RI-*Spl*II fragment derived from the plasmid pT796 and 0.05 µg of the phosphorylated synthetic DNA, obtained in the above, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained recombinant plasmid DNA solution, an *E. coli* HB101 was transformed to obtain the plasmid pT796H641L shown in Fig. 3. The thus obtained plasmid (10 µg) was allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to an electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence, and as a result, it was confirmed that a plasmid into which the DNA of interest was cloned was obtained.

Next, 3 µg of a plasmid pBluescript SK(-) (manufactured by Stratagene) was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 50 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 µg/ml BSA and the solution was further mixed with 10 units of a restriction enzyme *Xba*I (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 30 µl of a

buffer comprising 30 mM sodium acetate (pH 5.0), 100 mM sodium chloride, 1 mM zinc acetate and 5% glycerol, and further mixed with 30 units of a modification enzyme Mung Bean Nuclease (manufactured by Takara Shuzo) to carry out the reaction at 25°C for 15 minutes. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and further mixed with 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of a Blunt end-*ApaI* fragment of about 2.95 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 3 µg of the plasmid pKM641HF1 was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed with 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, and the 5' protruding end formed by the restriction enzyme digestion was changed to blunt end using DNA Blunting Kit (manufactured by Takara Shuzo). The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and further mixed with 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 0.2 µg of a Blunt end-*ApaI* fragment of about 0.44 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 0.1 μ g of Blunt end-ApaI fragment derived from the plasmid pBluescript SK(-) and 0.1 μ g of the Blunt end-ApaI fragment derived from the plasmid pKM641HF1, obtained in the above, were added to 20 μ l in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained recombinant plasmid DNA solution, an *E. coli* HB101 was transformed to obtain the plasmid pBS641H shown in Fig. 4.

Next, 3 μ g of the plasmid pT796H641L obtained in the above was added to 10 μ l of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT and the solution was further mixed with 10 units of a restriction enzyme ApaI (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 μ l of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT, 100 μ g/ml BSA and 0.01% Triton X-100, and further mixed with 10 units of a restriction enzyme NotI (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 0.2 μ g of an ApaI-NotI fragment of about 9.16 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 3 μ g of the plasmid pBS641H obtained in the above was added to 10 μ l of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT and the solution was further mixed with 10 units of a restriction enzyme ApaI (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 μ l of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT, 100 μ g/ml BSA and 0.01% Triton X-100, and further mixed with 10 units of a

restriction enzyme NotI (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an ApaI-NotI fragment of about 0.44 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 0.1 µg of the ApaI-NotI fragment derived from the plasmid pT796H641L and 0.1 µg of the ApaI-NotI fragment derived from the plasmid pBS641H, obtained in the above, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained recombinant plasmid DNA solution, an *E. coli* HB101 was transformed to obtain the plasmid pT641 shown in Fig. 5.

(2) Construction of an anti-GD3 CDR-grafted antibody KM871 transient expression vector

A transient expression vector for anti-GD3 CDR-grafted antibody was constructed as follows using the transient expression vector pT641 for anti-GD3 chimeric antibody described in the item 4(1) of Example 1 and the plasmids phKM641H and phKM641L described in the items 3(2) and(3) of Example 1.

First, 3 µg of the plasmid phKM641H obtained in the item 3(2) of Example 1 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT and the solution was further mixed with 10 units of a restriction enzyme ApaI (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT, 100 µg/ml BSA and 0.01% Triton X-100, and further mixed with 10 units of a restriction enzyme NotI (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour.

The reaction solution was fractionated by an agarose gel electrophoresis to recover about 0.2 µg of an ApaI-NotI fragment of about 0.44 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the

5 manufacture's instructions.

Next, 3 µg of the plasmid pT641 obtained in the item 4(1) of Example 1 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed

10 with 10 units of a restriction enzyme ApaI (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT,

15 100 µg/ml BSA and 0.01% Triton X-100, and further mixed with 10 units of a restriction enzyme NotI (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an ApaI-NotI

20 fragment of about 9.16 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 0.1 µg of the ApaI-NotI fragment derived from the plasmid pHKM641H and 0.1 µg of the ApaI-NotI fragment

25 derived from the plasmid pT641 were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained recombinant plasmid DNA solution, an *E. coli* HB101 was transformed to obtain the plasmid pT641HCDR shown in

30 Fig. 6.

Next, 3 µg of the plasmid pHKM641L obtained in the item 3(3) of Example 1 was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 µg/ml BSA, and

35 the solution was further mixed with 10 units of a

restriction enzyme *EcoRI* (manufactured by Takara Shuzo) and a restriction enzyme *SplI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 0.2 µg of an *EcoRI-SplI* fragment of about 0.39 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 3 µg of each of the plasmid pT641 obtained in the item 4(1) of Example 1 and the plasmid pT641HCDR obtained in the above was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 µg/ml BSA, and the solution was further mixed with 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) and a restriction enzyme *SplI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an *EcoRI-SplI* fragment of about 9.20 kb derived from each plasmid using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 0.1 µg of the *EcoRI-SplI* fragment derived from the plasmid pKM641L and 0.1 µg of the *EcoRI-SplI* fragment derived from the plasmid pT641, obtained in the above, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Also, 0.1 µg of the *EcoRI-SplI* fragment derived from the plasmid pKM641L and 0.1 µg of the *EcoRI-SplI* fragment derived from the plasmid pT641HCDR, obtained in the above, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using each of the thus obtained recombinant plasmid DNA solutions, an *E. coli* HB101 was transformed to obtain the plasmids pT641LCDR and pT641HLCDR shown in Fig. 7.

(3) Activity evaluation of anti-GD3 chimeric antibody and anti-GD3 CDR-grafted antibody by transient expression using animal cells

5 Transient expression of antibodies was carried out using the transient expression vector pT641 for anti-GD3 chimeric antibody, the transient expression vector pT641HLCDR for anti-GD3 CDR-grafted antibody and the transient expression vectors pT641HCDR and pT641LCDR for
10 anti-GD3 hybrid antibodies having V regions of mouse antibody and human CDR-grafted antibody, obtained in the items 4(1) and (2) of Example 1.

COS-7 cells (ATCC CRL 1651) were dispensed in 2 ml at a density of 1×10^5 cells/ml into a 6 well plate
15 (manufactured by Falcon) and cultured overnight at 37°C. By adding 2 µg of each expression vector to 100 µl of OPTI-MEM medium (manufactured by GIBCO BRL) and further adding a solution in which 10 µl of LIPOFECTAMINE Reagent (manufactured by GIBCO BRL) was added to 100 µl of the
20 OPTI-MEM medium, the reaction was carried out at room temperature for 40 minutes to effect formation of a DNA-liposome complex. The overnight-cultured COS-7 cells were washed twice with 2 ml of the OPTI-MEM medium (manufactured by GIBCO BRL), mixed with a solution prepared by adding 0.8
25 ml of OPTI-MEM medium to the complex-containing solution and cultured at 37°C for 7 hours, the solution was discarded and then the cells were mixed with 2 ml of 10% FBS-containing DME medium (manufactured by GIBCO BRL) and cultured at 37°C. After the introduction of each
30 expression vector, the culture supernatant was recovered 72 hours thereafter and its concentration operation was carried out as occasion demands, and then binding activity of the anti-GD3 humanized antibody for GD3 (manufactured by DIA-IATRON) in the culture supernatant was measured by the
35 ELISA method described in the item 2 of Example 1, by

measuring concentration of the anti-GD3 humanized antibody in the culture supernatant by the ELISA method described in the item 4 which will be described later and calculating the activity from the measured values as relative activity (%) when the activity of the positive control anti-GD3 chimeric antibody is defined as 100. Results are shown in Fig. 8. As shown in Fig. 8, the anti-GD3 hybrid antibody derived from the transient expression vector pT641HCDR (VH is derived from a human CDR-grafted antibody and VL is derived from a chimeric antibody) showed almost the same binding activity of that of the anti-GD3 hybrid antibody derived from the transient expression vector pT641, but the anti-GD3 hybrid antibody derived from the transient expression vector pT641LCDR (VH is derived from a chimeric antibody and VL is derived from a human CDR-grafted antibody) showed a binding activity of about 50% of the anti-GD3 chimeric antibody. In addition, the anti-GD3 CDR-grafted antibody derived from the transient expression vector pT641HLCADR showed a binding activity of only about 5% of the anti-GD3 chimeric antibody, showing considerable decrease in the activity.

Based on these results, binding activity of the anti-GD3 CDR-grafted antibody having VH and VL designed in the item 3(1) of Example 1 is considerably reduced and the main cause of the activity reduction is considered to be due to VL, but since about 50% of the activity is found by its combination with the VH of anti-GD3 chimeric antibody, it was suggested that there is a problem particularly at the interaction region with VH. Accordingly, an attempt was made to increase the antibody activity by further modifying amino acid residues of the VL designed in the item 3(1) of Example 1.

(4) Measurement of the concentration of humanized antibody in transient expression culture supernatant by ELISA method

A solution (50 μ l) prepared by diluting a goat anti-human IgG (γ -chain) antibody (manufactured by Medical & Biological Laboratories) 400 folds with PBS was dispensed in each well of a 96 well plate for ELISA use (manufactured by Greiner) and allowed to react at 4°C overnight. After discarding the antibody solution, 1% BSA-PBS was added in 100 μ l/well and allowed to react at room temperature for 1 hour to thereby block the remaining active groups. After discarding 1% BSA-PBS, the transient expression culture supernatants obtained in the item 4(3) of Example 1 or diluted solutions of the purified anti-GD3 chimeric antibody KM871 were added in 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction, each well was washed with Tween-PBS and then a solution prepared by diluting a peroxidase-labeled mouse anti-human κ L chain antibody (manufactured by Zymed) 500 folds with PBS was added in 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, an ABTS substrate solution (a solution prepared by dissolving 0.55 g of 2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid) in 1 L of 0.1 M citrate buffer (pH 4.2) and adding 1 μ l/ml of hydrogen peroxide just before the use) was added in 50 μ l/well for color development and OD415 was measured.

5. Increase of the binding activity by modifying amino acid residues of VL of anti-GD3 CDR-grafted antibody

Increase of the binding activity by modifying amino acid residues of VL of the anti-GD3 CDR-grafted antibody designed in the item 3(1) of Example 1 was carried out as follows.

(1) Modification of amino acid residues of VL of anti-GD3 CDR-grafted antibody and construction of cDNA encoding the modified VL

First, amino acid residues which are amino acid residues positioned in the interaction region of VH and VL and amino acid residues considered to be exerting influence on the three-dimensional structure of each CDR of VL and are also different from the amino acid residues of mouse antibody in the VL of human CDR-grafted antibody were identified from computer models of the three-dimensional structures of various antibodies constructed in the item 3(1) of Example 1. As a result, the 7th position Ser, the 8th position Pro, the 12th position Ser, the 41st position Gly, the 44th position Pro, the 72nd position Thr, the 77th position Ser, the 83rd position Phe and the 87th Tyr of hKM641L as the amino acid sequence of VL of anti-GD3 CDR-grafted antibody represented by SEQ ID NO:10 were identified. By replacing these amino acid residues by amino acid residues found in the mouse antibody, 8 VLs of modified anti-GD3 CDR-grafted antibody were designed. That is, the 8th position Pro, the 12th position Ser, the 44th position Pro and the 87th position Tyr among the hKM641L amino sequence were respectively replaced by Ala, Pro, Val and Phe in the case of hKM641NL, and the 7th position Ser, the 8th position Pro and the 12th position Ser among the hKM641L amino sequence respectively by Thr, Ala and Pro in the case of hKM641Lm-1, the 87th position Tyr among the hKM641L amino sequence by Phe in the case of hKM641Lm-4, the 41st position Gly and the 44th position Pro among the hKM641L amino sequence respectively by Asp and Val in the case of hKM641Lm-6, the 72nd position Thr, the 77th position Ser and the 83rd position Phe among the hKM641L amino sequence respectively by Ser, Asn and Ile in the case of hKM641Lm-7, the 77th position Ser among the hKM641L amino sequence by Asn in the case of hKM641Lm-8, the 83rd

position Phe and the 87th position Tyr among the hKM641L amino sequence respectively by Ile and Phe in the case of hKM641Lm-9, and the 41st position Gly, the 44th position Pro and the 83rd position Phe among the hKM641L amino sequence respectively by Asp, Val and Ile in the case of hKM641Lm-69. Among these modified VLs, cDNA encoding each of 6 modified VLs excluding hKM641NL and hKM641Lm-69 was constructed as follows by PCR-aided mutagenesis. That is, antisense chain and sense chain DNA primers were synthesized using an automatic DNA synthesizer (380A, manufactured by Applied Biosystems) for introducing mutation, and a first PCR was carried out in accordance with the method described in the item 3(2) of Example 1 using 1 ng of the plasmid phKM641L as the template and 0.5 μ M in final concentration of M13 primer RV (manufactured by Takara Shuzo) and the antisense chain DNA primer and M13 primer M4 (manufactured by Takara Shuzo) and the sense chain DNA primer. Each of the reaction solutions was purified using QIAquick PCR Purification Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions by eluting with 20 μ l of 10 mM Tris-HCl (pH 8.0), and then a second PCR was carried out using 5 μ l of each eluate in accordance with the method described in the item 3(2) of Example 1. The reaction solution was purified using QIAquick PCR Purification Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions and made into 30 μ l of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 μ g/ml BSA, and 10 units of a restriction enzyme *Eco*RI (manufactured by Takara Shuzo) and a restriction enzyme *Spl*I (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 0.2 μ g of an *Eco*RI-*Spl*I fragment of about 0.39 kb using QIAquick Gel

Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

5 Next, 0.1 µg of the *EcoRI*-*SplI* fragment of the PCR product of VL of the modified anti-GD3 CDR-grafted antibody obtained in the above and 0.1 µg of the *EcoRI*-*SplI* fragment of the plasmid pBSL3 obtained in the item 3(3) of Example 1 were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained recombinant plasmid 10 DNA solution, an *E. coli* HB101 was transformed. Each plasmid DNA was produced from 10 clones of the transformants, allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to 15 electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence to thereby obtain a plasmid having cDNA to which the intended modification was applied.

20 Specifically, a plasmid phKM641Lm-1 having the nucleotide sequence represented by SEQ ID NO:27 was obtained by carrying out a series of the above operations using the synthetic DNA of SEQ ID NO:25 as the antisense chain DNA primer and the synthetic DNA of SEQ ID NO:26 as the sense chain DNA primer. The hKM641Lm-1 as an amino 25 acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:27.

A plasmid phKM641Lm-4 having the nucleotide sequence represented by SEQ ID NO:30 was obtained by carrying out a series of the above operations using the 30 synthetic DNA of SEQ ID NO:28 as the antisense chain DNA primer and the synthetic DNA of SEQ ID NO:29 as the sense chain DNA primer. The hKM641Lm-4 as an amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:30.

A plasmid pHKM641Lm-6 having the nucleotide sequence represented by SEQ ID NO:33 was obtained by carrying out a series of the above operations using the synthetic DNA of SEQ ID NO:31 as the antisense chain DNA primer and the synthetic DNA of SEQ ID NO:32 as the sense chain DNA primer. The hKM641Lm-6 as an amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:33.

A plasmid pHKM641Lm-7 having the nucleotide sequence represented by SEQ ID NO:36 was obtained by carrying out a series of the above operations using the synthetic DNA of SEQ ID NO:34 as the antisense chain DNA primer and the synthetic DNA of SEQ ID NO:35 as the sense chain DNA primer. The hKM641Lm-7 as an amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:36.

A plasmid pHKM641Lm-8 having the nucleotide sequence represented by SEQ ID NO:39 was obtained by carrying out a series of the above operations using the synthetic DNA of SEQ ID NO:37 as the antisense chain DNA primer and the synthetic DNA of SEQ ID NO:38 as the sense chain DNA primer. The hKM641Lm-8 as an amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:39.

A plasmid pHKM641Lm-9 having the nucleotide sequence represented by SEQ ID NO:42 was obtained by carrying out a series of the above operations using the synthetic DNA of SEQ ID NO:40 as the antisense chain DNA primer and the synthetic DNA of SEQ ID NO:41 as the sense chain DNA primer. The hKM641Lm-9 as an amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:42.

Among the modified VLs, cDNA encoding the hKM641NL was constructed by synthesizing 6 synthetic DNA fragments of SEQ ID NOS:17, 22 and 43 to 46 using an automatic DNA

synthesizer (380A, manufactured by Applied Biosystems) and carrying out the procedure described in the item 3(3) of Example 1 using these fragments. As a result, a plasmid pHKM641NL having the nucleotide sequence represented by SEQ ID NO:47 in which the modification of interest was effected was obtained. The hKM641NL as an amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:47.

Among the modified VLs, cDNA encoding the hKM641Lm-69 was constructed as follows using the plasmids pHKM641Lm-6 and pHKM641Lm-9 having cDNA of modified VL obtained in the above.

First, 3 μ g of the plasmid pHKM641Lm-6 was added to 10 μ l of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM of sodium chloride, 10 mM magnesium chloride and 1 mM DTT and 10 units of restriction enzymes *Eco*RI (manufactured by Takara Shuzo) and *Pst*I (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 0.3 μ g of an *Eco*RI-*Pst*I fragment of about 0.30 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 3 μ g of the plasmid pHKM641Lm-9 was added to 10 μ l of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM of sodium chloride, 10 mM magnesium chloride and 1 mM DTT and 10 units of restriction enzymes *Eco*RI (manufactured by Takara Shuzo) and *Pst*I (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 2 μ g of an *Eco*RI-*Pst*I fragment of about 3.05 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 0.1 µg of the EcoRI-PstI fragment derived from the plasmid phKM641Lm-6 and 0.1 µg of the EcoRI-PstI fragment derived from the plasmid phKM641Lm-9, both obtained in the above, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained recombinant plasmid DNA solution, an *E. coli* HB101 was transformed to obtain the plasmid phKM641Lm-69 shown in Fig. 9. The reaction was carried out using 10 µg of the thus obtained plasmid in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then the resulting mixture was subjected to electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence, and as a result, it was confirmed that it has the nucleotide sequence represented by SEQ ID NO:48 in which the modification of interest was carried out. The hKM641Lm-69 as an amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:48.

(2) Construction of transient expression vector for anti-GD3 CDR-grafted antibodies having modified VL

Transient expression vectors for anti-GD3 CDR-grafted antibodies having various modified VLs were constructed as follows using the plasmids having cDNA molecules encoding various modified VLs, obtained in the item 5(1) of Example 1, and the transient expression vector pT641HCDCR for anti-GD3 hybrid antibodies obtained in the item 4(2) of Example 1.

First, 3 µg of each of the plasmids phKM641NL, phKM641Lm-1, phKM641Lm-4, phKM641Lm-6, phKM641Lm-7, phKM641Lm-8, phKM641Lm-9 and phKM641Lm-69 obtained in the item 5(1) of Example 1 was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM of sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 µg/ml

of BSA, and 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) and a restriction enzyme *SplI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 0.2 µg of an *EcoRI-SplI* fragment of about 0.39 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 0.1 µg of each of the thus obtained *EcoRI-SplI* fragments of various modified VLs and 0.1 µg of the *EcoRI-SplI* fragment of the transient expression vector pT641HCDR for anti-GD3 hybrid antibody obtained in the item 4(2) of Example 1, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained respective recombinant plasmid DNA solutions, an *E. coli* HB101 was transformed to obtain the transient expression vectors for anti-GD3 CDR-grafted antibodies having various modified VLs, pT641HLCDRNL, pT641HLCDRM-1, pT641HLCDRM-4, pT641HLCDRM-6, pT641HLCDRM-7, pT641HLCDRM-8, pT641HLCDRM-9 and pT641HLCDRM-69, shown in Fig. 10.

(3) Evaluation of activity of anti-GD3 CDR-grafted antibody having modified VLs by transient expression using animal cell

Transient expression and evaluation of activity of each antibody were carried out in accordance with the method described in the item 4(3) of Example 1 using the transient expression vector pT641 for anti-GD3 chimeric antibody and the transient expression vector pT641HLCDR for anti-GD3 CDR-grafted antibody, both obtained in the item 4(2) of Example 1, and the transient expression vectors, pT641HLCDRNL, pT641HLCDRM-1, pT641HLCDRM-4, pT641HLCDRM-6, pT641HLCDRM-7, pT641HLCDRM-8, pT641HLCDRM-9 and

pT641HLCDRM-69, for anti-GD3 CDR-grafted antibodies having various modified VLs, obtained in the item 5(2) of Example 1. Results are shown in Fig. 11. As shown in Fig. 11, the modified anti-GD3 CDR-grafted antibodies derived from the transient expression vectors, pT641HLCDRM-6, pT641HLCDRM-7, pT641HLCDRM-9 and pT641HLCDRM-69, showed increased binding activity in comparison with the anti-GD3 CDR-grafted antibodies before modification, particularly, the modified anti-GD3 CDR-grafted antibody derived from the transient expression vector pT641HLCDRM-69 showed its binding activity of about 80% of the anti-GD3 chimeric antibody. On the other hand, the modified anti-GD3 CDR-grafted antibodies derived from other transient expression vectors showed no increase in the binding activity in comparison with the anti-GD3 CDR-grafted antibodies before modification. Based on these results, it was found that, among the modified amino acid residues of VL identified in the item 5(1) of Example 1, the 41st position, 44th position, 83rd position and 87th position amino acid residues greatly contribute to the increase of binding activity. It was also found that simultaneous substitution of two amino acid residues of the 41st position and 44th position or the 83rd position and 87th position contributes to synergistic increase in the activity, and simultaneous modification of four amino acid residues of the 41st position, 44th position, 83rd position and 87th position also contributes to synergistic increase in the activity. Based on the computer models of three-dimensional structures of V regions of various antibodies constructed in the item 3(1) of Example 1, it was suggested that the 41st position and 44th position amino acid residues exert influence on the interaction with VH, and the 83rd position and 87th position amino acid residues on the three-dimensional structure of CDR3 of VL, and it was considered that three-dimensional structure of an antibody as a whole

is suitably maintained by modifying these amino acid residues into amino acid residues found in a mouse antibody and, as a result, its binding activity increases. This information shows that it is necessary to take the interaction regions of VH and VL also into consideration in preparing human CDR-grafted antibodies and that it is necessary to carry out various examinations on the identification of these interaction regions based on the information from three-dimensional structures of antibody V regions. With regard to these interaction regions, it can be easily analogize that they are different in individual antibodies and it is difficult to find a general law at present, so that it is necessary to carry out trial and error in response to each antibody of interest.

6. Stable expression of anti-GD3 CDR-grafted antibody using animal cell

Based on the results described in the item 5(3) of Example 1, it was suggested that the anti-GD3 CDR-grafted antibodies derived from transient expression vectors pT641HLCDRIm-9 and pT641HLCDRIm-69 have about 20% and about 80% of binding activity for GD3, respectively, in comparison with anti-GD3 chimeric antibodies. Accordingly, in order to evaluate the activity of these anti-GD3 CDR-grafted antibodies more in detail, transformed cell lines capable of stably expressing anti-GD3 CDR-grafted antibodies were obtained by the following method, and purification of anti-GD3 CDR-grafted antibodies from culture supernatants of the transformed cell lines was carried out. Also, for comparison, stable expression and purification of the anti-GD3 CDR-grafted antibody derived from the transient expression vector pT641HLCDR were carried out in the same manner.

(1) Construction of stable expression vector for anti-GD3 CDR-grafted antibody

Stable expression vectors were constructed by introducing a resistance gene against an agent G418 and the dhfr gene into various transient expression vectors constructed in the item 5(2) of Example 1.

First, 3 µg of the humanized antibody expression vector pKANTEX93 described in WO 97/10354 was added to 10 µl of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and 10 units of a restriction enzyme BamHI (manufactured by Takara Shuzo) and a restriction enzyme XhoI (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 1 µg of a BamHI-XhoI fragment of about 8.68 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 3 µg of each of the transient expression vectors, pT641HLCDRIm-9, pT641HLCDRIm-69 and pT641HLCDR, was added to 10 µl of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and 10 units of a restriction enzyme BamHI (manufactured by Takara Shuzo), a restriction enzyme XhoI (manufactured by Takara Shuzo) and a restriction enzyme StuI (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 1 µg of a BamHI-XhoI fragment of about 4.90 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 0.1 µg of the BamHI-XhoI fragment derived from the plasmid pKANTEX93 and 0.1 µg of the BamHI-XhoI fragment derived from each of the transient expression

vectors, obtained in the above, were added to 20 μ l in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using each of the thus obtained recombinant plasmid DNA solutions, an *E. coli* HB101 was transformed to obtain the anti-GD3 CDR-grafted antibody stable expression vectors, pKANTEX641HLCDRM-9, pKANTEX641HLCDRM-69 and pKANTEX641HLCDR, shown in Fig. 12.

10 (2) Stable expression of anti-GD3 CDR-grafted antibody using animal cell

Using the various stable expression vectors for anti-GD3 CDR-grafted antibody obtained in the item 6(1) of Example 1, expression of anti-GD3 CDR-grafted antibody in animal cells was carried out as follows.

Each of the anti-GD3 CDR-grafted antibody expression vectors (4 μ g) was introduced into 4×10^5 cells of a rat myeloma cell line YB2/0 cell (ATCC CRL 1662) by the electroporation method (Cytotechnology, 2, 133 (1990)), and the cells were suspended in 40 ml of RPMI1640-FBS(19) (RPMI1640 medium containing 10% of fetal bovine serum (FBS)) and dispensed in 200 μ l/well into a 96 well microtiter plate (manufactured by Sumitomo Bakelite). After culturing at 37°C for 24 hours in a 5% CO₂ incubator, G418 was added to give a concentration of 0.5 mg/ml, followed by culturing for 1 to 2 weeks. Culture supernatants were recovered from wells in which colonies of transformants showing G418-resistance were formed and became confluent, and the antigen binding activity of various anti-GD3 CDR-grafted antibodies in the supernatants was measured by the ELISA shown in the item 2 of Example 1 (a peroxidase-labeled goat anti-human IgG(γ) antibody was used as the secondary antibody).

With regard to the transformants of wells in which expression of anti-GD3 CDR-grafted antibodies was found in

the culture supernatants, in order to increase antibody expression quantity making use of a dhfr gene amplification system, they were suspended in the RPMI1640-FBS(10) medium containing 0.5 mg/ml of G418 and 50 nM methotrexate (hereinafter referred to as "MTX": manufactured by SIGMA) as an inhibitor of the dhfr gene product dihydrofolate reductase (hereinafter referred to as "DHFR") to give a density of 1 to 2×10^5 cells/ml, and dispensed in 2 ml into a 24 well plate (manufactured by Greiner). By culturing at 37°C for 1 to 2 weeks in a 5% CO₂ incubator, transformants showing 50 nM MTX resistance were induced. When the transformants became confluent, the antigen binding activity of various anti-GD3 CDR-grafted antibodies in the supernatants was measured by the ELISA shown in the item 2 of Example 1. With regard to the transformants of wells in which expression of anti-GD3 CDR-grafted antibodies was found in the culture supernatants, the MTX concentration was increased to 100 nM and then to 200 nM by the method similar to the above, and transformants capable of growing in the RPMI1640-FBS(10) medium containing 0.5 mg/ml of G418 and 200 nM of MTX and of highly expressing anti-GD3 CDR-grafted antibodies were finally obtained. With regard to the thus obtained transformants, single cell isolation (cloning) was carried out twice by limiting dilution analysis. The transformed cell clones obtained by introducing the stable expression vectors, pKANTEX641HLCDR_{Lm}-9, pKANTEX641HLCDR_{Lm}-69 and pKANTEX641HLCDR, were named KM8870, KM8871 and KM8869, respectively, and expression levels of anti-GD3 CDR-grafted antibody by transformed cell clones were about 5 µg/10⁶ cells/24 hr, about 10 µg/10⁶ cells/24 hr and about 30 µg/10⁶ cells/24 hr, respectively.

(3) Purification of anti-GD3 CDR-grafted antibody from culture filtrate

Each of the transformed cell clones, KM8870, KM8871 and KM8869, obtained in the item 6(2) of Example 1 which
5 express various anti-GD3 CDR-grafted antibodies was suspended in GIT medium (manufactured by Nippon Pharmaceutical) containing 200 nM MTX to give a density of 1 to 2×10^5 cells/ml, and dispensed in 200 ml into 175 cm² flasks (manufactured by Greiner). The cells were cultured
10 at 37°C for 5 to 7 days until they became confluent and then the culture supernatant was recovered. Each anti-GD3 CDR-grafted antibody was purified from the culture supernatant using Prosep-A (manufactured by Bioprocessing) column in accordance with the manufacture's instructions.
15 About 3 mg of anti-GD3 CDR-grafted antibody KM8870 was obtained from 500 ml of the culture supernatant of KM8870, and about 25 mg of anti-GD3 CDR-grafted antibody KM8871 from 1,600 ml of the culture supernatant of KM8871 and about 65 mg of anti-GD3 CDR-grafted antibody KM8869 from
20 1,000 ml of the culture supernatant of KM8869. About 4 µg of each of the thus obtained anti-GD3 CDR-grafted antibodies was subjected to an electrophoresis in accordance with the known method (*Nature*, 227, 680 (1970)) to examine its molecular weight and purification degree.
25 Results are shown in Fig. 13. As shown in Fig. 13, molecular weight of each purified anti-GD3 CDR-grafted antibody was about 150 kilodaltons (hereinafter referred to as "Kd") under non-reducing conditions, and two bands of about 50 Kd and about 25 Kd were found under reducing
30 conditions. The molecular weights almost coincided with the molecular weights deduced from the cDNA nucleotide sequences of H chain and L chain of the antibody (H chain: about 49 Kd, L chain: about 24 Kd, whole molecule: about 146 Kd), and also coincided with the reports that the IgG
35 type antibody has a molecular weight of about 150 Kd under

non-reducing conditions and is degraded into H chains having a molecular weight of about 50 Kd and L chains having a molecular weight of about 25 Kd under reducing conditions due to cutting of the disulfide bond (hereinafter referred to as "S-S bond") in the molecule (Antibodies: A Laboratory Manual, Monoclonal Antibodies: Principles and Practice), so that it was confirmed that each anti-GD3 CDR-grafted antibody was expressed as an antibody molecule of a correct structure. Also, when N-terminal amino acid sequences of the H chain and L chain of each of the purified anti-GD3 CDR-grafted antibodies were analyzed by Edman degradation using a protein sequencer (470A, manufactured by Applied Biosystems), it was confirmed that they coincide with the H chain and L chain N-terminal amino acid sequences deduced from respective cDNA nucleotide sequences.

7. Evaluation of activity of anti-GD3 CDR-grafted antibody

(1) Reactivity of anti-GD3 CDR-grafted antibody with GD3 (ELISA)

Reactivity of purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 with GD3 (manufactured by DIA-IATRON) was measured by the ELISA shown in the item 2 of Example 1. Fig. 14 shows a result of the examination of the reactivity carried out by fixing the amount of GD3 to be adsorbed to each well of a plate for ELISA to 20 pmol/well and changing the concentration of anti-GD3 chimeric antibody KM871 and anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 to be added. As shown in Fig. 14, among the anti-GD3 CDR-grafted antibodies, KM8871 showed the most highest binding activity which was about 1/2 of the anti-GD3 chimeric antibody KM871. Fig. 15 shows a result of the examination of the reactivity of a constant concentration (10 µg/ml) of anti-GD3 chimeric antibody KM871 and anti-GD3 CDR-grafted antibodies KM8869, KM8870

and KM8871, carried out by changing the amount of GD3 to be adsorbed to each well of a plate for ELISA use. As shown in Fig. 15, among the anti-GD3 CDR-grafted antibodies, KM8871 showed the most highest binding activity which was equivalent to the anti-GD3 chimeric antibody KM871. Fig. 16 shows a result of the examination of the reactivity of a constant concentration (10 µg/ml) of anti-GD3 chimeric antibody KM871 and anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871, carried out by changing the kind of ganglioside to be adsorbed to each well of a plate for ELISA use (adsorption amount: 20 pmol/well). A total of 11 kinds of ganglioside, GM1, N-acetylGM2 (manufactured by Boehringer Mannheim, hereinafter referred to as "AcGM2"), N-glycolylGM2 (hereinafter referred to as "GcGM2"), N-acetylGM3 (hereinafter referred to as "AcGM3"), N-glycolylGM3 (hereinafter referred to as "GcGM3"), GD1a and GD1b (manufactured by DIA-IATRON), GD2 and GD3 (manufactured by DIA-IATRON), Q1b (manufactured by DIA-IATRON) and GT1b (manufactured by Funakoshi), were used. In this case, GM1 and GD1a were purified from bovine brain, and N-glycolylGM2 and N-glycolylGM3 from mouse liver, N-acetylGM3 from canine erythrocyte and GD2 from a human neuroblastoma culture cell line IMR32 (ATCC CCL 127), respectively in accordance with the known method (*J. Biol. Chem.*, 263, 10915 (1988)). As shown in Fig. 16, each of the anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 bound strongly to GD3 and weakly to Q1b similar to the case of the anti-GD3 chimeric antibody KM871 but did not bind to other gangliosides. Thus, it was shown that they maintain the binding specificity of chimeric antibody KM871.

(2) Reactivity of anti-GD3 CDR-grafted antibody with human cancer cell (immunofluorescent method)

Reactivity of purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 with human cancer cells was measured as follows. That is, 1×10^6 cells of each of human melanoma culture cell lines G-361 (ATCC CRL 1424) and SK-MEL-28 (ATCC HTB72) were suspended in PBS, put into microtubes and centrifuged (2,000 rpm for 2 minutes), and the thus washed cells were stirred after adding 50 μ l of the anti-GD3 chimeric antibody KM871 or the anti-GD3 CDR-grafted antibody, KM8869, KM8870 or KM8871 (a solution adjusted to 5 μ g/ml with 1% BSA-PBS) and then allowed to react at 4°C for 1 hour. After the reaction, followed by centrifugation three times with PBS for washing, 20 μ l of rabbit anti-human IgG (H + L) F(ab')₂ solution (manufactured by Wako Pure chemical Industries, used by diluting 30 times with 1% BSA-PBS) fluorescence-labeled with fluorescein isothiocyanate (hereinafter referred to as "FITC") was added thereto, followed by stirring, and then the reaction was carried out at 4°C for 1 hour. After the reaction, followed by centrifugation three times with PBS for washing, the cells were again suspended in PBS to carry out the analysis using a flow cytometer EPICS Elite (manufactured by COULTER). As a control, the analysis was carried out by the same procedure without adding antibodies. Results are shown in Fig. 17. As shown in Fig. 17, all of the anti-GD3 chimeric antibody KM871 and the anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 showed the reactivity with both cell lines. Among the anti-GD3 CDR-grafted antibodies, KM8871 showed the most strongest reactivity. The above results show that the anti-GD3 CDR-grafted antibodies are useful in the diagnosis, treatment and the like of GD3-positive human tumors including melanoma.

(3) *In vitro* cytotoxic activity of anti-GD3 CDR-grafted antibodies (CDC activity)

In order to evaluate *in vitro* cytotoxic activity of the purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871, the CDC activity was measured by the following method.

a. Preparation of target cell suspension

Each of human melanoma culture cell lines, G-361 (ATCC CRL 1424) and SK-MEL-28 (ATCC HTB72), was cultured in RPMI1640-FBS(10) medium and adjusted to 5×10^6 cells, and the cells were isotope-labeled by adding 3.7 MBq equivalent of a radioactive substance $\text{Na}_2^{51}\text{CrO}_4$ and carrying out the reaction at 37°C for 1 hour. After the reaction, the cells were washed three times by their suspension in RPMI1640-FBS(10) medium and centrifugation, re-suspended in the medium and then incubated in ice at 4°C for 30 minutes to thereby spontaneously release the radioactive substance. After centrifugation, the precipitate was adjusted to 1×10^6 cells/ml by adding 5 ml of RPMI1640-FBS(10) medium and used as the target cell suspension.

b. Preparation of complement solution

Sera of three healthy persons were mixed and used as the human complement source. At the time of its use, it was diluted to 15% vol/vol with RPMI1640-FBS(10) medium and used as the complement solution.

c. Measurement of CDC activity

To each well of a 96 well U bottom plate (manufactured by Falcon), 50 μl of the target cell suspension prepared in "a" was added (5×10^4 cells/well), and then the anti-GD3 chimeric antibody KM871 or the anti-GD3 CDR-grafted antibody, KM8869, KM8870 or KM8871, was added to give a final concentration of 0.05 to 50 $\mu\text{g/ml}$ and

allowed to react at room temperature for 30 minutes. After
 the reaction, the plate was centrifuged to remove the
 supernatant, and 150 μ l of the human complement solution
 prepared in "b" was added thereto and allowed to react at
 5 37°C for 1 hour. After centrifugation, the amount of ^{51}Cr
 released in the supernatant was measured by a γ -counter.
 The amount of the spontaneously dissociated ^{51}Cr was
 calculated by carrying out the same procedure using the
 medium alone instead of the antibody solution and
 10 complement solution and measuring the amount of ^{51}Cr in the
 supernatant. Amount of the total dissociated ^{51}Cr was
 calculated by carrying out the same procedure by adding the
 medium alone instead of the antibody solution, and 5 N
 sodium hydroxide instead of the complement solution, and
 15 measuring the amount of ^{51}Cr in the supernatant. The CDC
 activity was calculated by the following equation.

$$\begin{aligned}
 & \text{(amount of } ^{51}\text{Cr in sample supernatant)} \\
 & - \text{(amount of spontaneously released } ^{51}\text{Cr)} \\
 20 \quad \text{CDC activity (\%)} = & \frac{\quad}{\text{(amount of total } ^{51}\text{Cr)} - \text{(amount of spontaneously released } ^{51}\text{Cr)}} \times 100
 \end{aligned}$$

Results are shown in Fig. 18. As shown in Fig. 18,
 25 it was found that all of the anti-GD3 chimeric antibody
 KM871 and the anti-GD3 CDR-grafted antibodies KM8869,
 KM8870 and KM8871 show the CDC activity for both cell lines,
 particularly show remarkably strong cytotoxic activity for
 the human melanoma culture cell line G-361. Among the
 30 anti-GD3 CDR-grafted antibodies, KM8871 showed the most
 stronger cytotoxic activity which was about 1/3 of the
 anti-GD3 chimeric antibody KM871.

(4) *In vitro* cytotoxic activity of anti-GD3 CDR-grafted antibody (ADCC activity)

In order to evaluate *in vitro* cytotoxic activity of the purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871, the ADCC activity was measured by the following method.

a. Preparation of target cell suspension

Each of human melanoma culture cell lines, G-361 (ATCC CRL 1424) and SK-MEL-28 (ATCC HTB72), was cultured in RPMI1640-FBS(10) medium and adjusted to 1×10^6 cells, and the cells were isotope-labeled by adding 1.85 MBq equivalent of a radioactive substance $\text{Na}_2^{51}\text{CrO}_4$ and carrying out the reaction at 37°C for 1 hour. After the reaction, the cells were washed three times by their suspension in RPMI1640-FBS(10) medium and centrifugation, re-suspended in the medium and then incubated in ice at 4°C for 30 minutes to thereby spontaneously release the radioactive substance. After centrifugation, the precipitate was adjusted to 2×10^5 cells/ml by adding 5 ml of RPMI1640-FBS(10) medium and used as the target cell suspension.

b. Preparation of effector cell suspension

Healthy human vein blood (50 ml) was collected, and 0.5 ml of heparin sodium (manufactured by Takeda Pharmaceutical) was gently added thereto. The resulting mixture was centrifuged ($1,500$ to $1,800 \times g$ for 30 minutes) using Polymorphprep (manufactured by Nycomed Pharma AS) in accordance with the manufacture's instructions to separate a mononuclear cell layer. The separated cells were three times centrifuged ($1,500$ to $1,800 \times g$ for 5 minutes) with RPMI1640-FBS(10) medium for washing and then re-suspended in the medium to give a density of 5×10^6 cells/ml to be used as the effector cell suspension.

c. Measurement of ADCC activity

Into each well of a 96 well U bottom plate (manufactured by Falcon), 50 μ l of the target cell suspension prepared in "a" was dispensed (1×10^4 cells/well).

Next, 100 μ l of the effector cell suspension prepared in "b" was added (5×10^5 cells/well, ratio of the effector cells to the target cells becomes 50:1). Thereafter, each of the anti-GD3 chimeric antibody KM871 and the anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 was added to give a final concentration of 0.05 to 50 μ g/ml and allowed to react at 37°C for 4 hours. After the reaction, the plate was centrifuged and the amount of ^{51}Cr in the supernatant was measured by a γ -counter. The amount of the spontaneously dissociated ^{51}Cr was calculated by carrying out the same procedure using the medium alone instead of the effector cell suspension and antibody solution and measuring the amount of ^{51}Cr in the supernatant. The amount of the total dissociated ^{51}Cr was calculated by carrying out the same procedure by adding the medium alone instead of the antibody solution, and 5 N sodium hydroxide instead of the effector cell suspension, and measuring the amount of ^{51}Cr in the supernatant. The ADCC activity was calculated by the following equation.

$$\text{ADCC activity (\%)} = \frac{(\text{amount of } ^{51}\text{Cr in sample supernatant}) - (\text{amount of spontaneously released } ^{51}\text{Cr})}{(\text{amount of total } ^{51}\text{Cr}) - (\text{amount of spontaneously released } ^{51}\text{Cr})} \times 100$$

Results are shown in Fig. 19. As shown in Fig. 19, it was found that all of the anti-GD3 chimeric antibody KM871 and the anti-GD3 CDR-grafted antibodies, KM8869, KM8870 and KM8871, show strong ADCC activity for both cell lines. Among the anti-GD3 chimeric antibody KM871 and the

anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871, difference in the cytotoxic activity was not found.

The above results show that the anti-GD3 CDR-grafted antibodies are useful in the diagnosis, treatment and the like of GD3-positive human tumors similar to the case of the anti-GD3 chimeric antibody KM871. In addition, it is expected that immunogenicity of the anti-GD3 CDR-grafted antibodies is reduced in human and the effect is further prolonged.

The transformed cell clone KM8871 capable of producing KM8871 which showed the most high activity among the anti-GD3 CDR-grafted antibodies has been deposited on July 22, 1999, as FERM BP-6790 in National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology (Higashi 1-1-3, Tsukuba, Ibaraki, Japan).

Example 2

Preparation of fusion protein of anti-GD3 CDR-grafted antibody with human cytokine:

As an example of a fusion protein of anti-GD3 CDR-grafted antibody with human cytokine, a fusion protein of the anti-GD3 CDR-grafted antibody KM8871 with human IL-2, KM8871-hIL-2, was prepared as follows and evaluation of the activity was carried out.

1. Construction of cDNA encoding a fusion protein of hCyl with hIL-2

(1) Construction of a plasmid pBSΔhCyl-IL-2 having a cDNA comprised of about 65 bases of 3'-terminal of hCyl cDNA and full length cDNA of matured type hIL-2

A plasmid pBluescript SK(-) (3 μg, manufactured by Stratagene) was added to 10 μl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed with 10 units of a

restriction enzyme *ApaI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and further mixed with 10 units of a restriction enzyme *BamHI* (manufactured by Takara Shuzo) to carry out the reaction at 30°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an *ApaI-BamHI* fragment of about 2.95 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, synthetic DNA fragments respectively having the nucleotide sequences represented by SEQ ID NO:49 and SEQ ID NO:50 were synthesized using an automatic DNA synthesizer (380A, manufactured by Applied Biosystems). A 0.3 µg of each of the thus obtained synthetic DNA fragments was added to 15 µl of sterile water and heated at 65°C for 5 minutes. After allowing the reaction solution to stand at room temperature for 30 minutes, this was mixed with 2 µl of a 10 × buffer solution (500 mM Tris-HCl (pH 7.6), 100 mM magnesium chloride, 50 mM DTT) and 2 µl of 10 mM ATP and further mixed with 10 units of T4 Polynucleotide Kinase (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 30 minutes to thereby phosphorylate the 5'-terminal.

Next, 0.1 µg of the *ApaI-BamHI* fragment derived from the plasmid pBluescript SK(-) and 0.05 µg of the phosphorylated synthetic DNA, both obtained in the above, were added to 10 µl in total volume of sterile water and ligated using DNA Ligation Kit Ver. 2 (manufactured by Takara Shuzo) in accordance with the manufacture's instructions. Using the thus obtained recombinant plasmid DNA solution, an *E. coli* line DH5α (manufactured by Stratagene) was transformed to obtain the plasmid pBSA-B

shown in Fig. 20. The thus obtained plasmid (10 µg) was allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to an electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence, and as a result, it was confirmed that a plasmid into which the DNA of interest was cloned was obtained.

Next, 3 µg of the thus obtained plasmid pBSA-B was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed with 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 33 mM Tris-acetate (pH 7.9), 66 mM potassium acetate, 10 mM magnesium acetate, 0.5 mM DTT and 100 µg/ml BSA, and further mixed with 10 units of a restriction enzyme *SmaI* (manufactured by Takara Shuzo) to carry out the reaction at 30°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an *EcoRI-SmaI* fragment of about 3.00 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, the following PCR was carried out using a plasmid pILL4 containing full length cDNA of matured type hIL-2 (*Agric. Biol. Chem.*, 51, 1135 (1987)) as the template. The plasmid pILL4 (1 ng) was added to 100 µl of a reaction solution (1 × concentration Ex Taq buffer (manufactured by Takara Shuzo), 200 µM dNTPs, 1.0 µM rev1 primer (SEQ ID NO:51), 1.0 µM fw2 primer (SEQ ID NO:52) and 2.5 units of TaKaRa Ex Taq DNA polymerase (manufactured by Takara Shuzo)), and the solution was covered with 100 µl of mineral oil and set to a DNA thermal cycler (PJ480, manufactured by PERKIN ELMER) to carry out the reaction at

94°C for 3 minutes, subsequent 30 cycles of each cycle at 96°C for 30 seconds, 55°C for 1 minute and 72°C for 1 minute, and finally at 72°C for 7 minutes. The reaction solution was precipitated with ethanol, added to 30 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride and 1 mM DTT, and further mixed with 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 33 mM Tris-acetate (pH 7.9), 66 mM potassium acetate, 10 mM magnesium acetate, 0.5 mM DTT and 100 µg/ml BSA, and further mixed with 10 units of a restriction enzyme *SmaI* (manufactured by Takara Shuzo) to carry out the reaction at 30°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 1 µg of an *EcoRI-SmaI* fragment of about 0.41 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 0.1 µg of the thus obtained *EcoRI-SmaI* fragment of the plasmid pBSA-B and 0.1 µg of the *EcoRI-SmaI* fragment of the PCR product of full length cDNA of matured type hIL-2 were added to 10 µl in total volume of sterile water and ligated using DNA Ligation Kit Ver. 2 (manufactured by Takara Shuzo) in accordance with the manufacture's instructions. Using the thus obtained recombinant plasmid DNA solution, an *E. coli* DH5α was transformed. Each plasmid DNA was prepared from 10 clones of the transformants, allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to an electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine nucleotide sequence of the inserted cDNA, and as a result, the plasmid

pBSΔhCyl-IL-2 shown in Fig. 21 having the nucleotide sequence of interest was obtained.

(2) Construction of a plasmid pBShCyl-IL-2 having a cDNA comprised of hCyl full length cDNA and full length cDNA of matured type hIL-2

First, 3 µg of the plasmid pBShCyl described in Japanese Published Unexamined Patent Application No. 257893/98 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed with 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and further mixed with 10 units of a restriction enzyme *EcoT22I* (manufactured by Takara Shuzo) to carry out the reaction at 30°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 1 µg of an *ApaI*-*EcoT22I* fragment of about 0.92 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 3 µg of the plasmid pBSΔhCyl-IL-2 obtained in the item 1(1) of Example 2 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed with 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM sodium chloride, and further mixed with 10 units of a restriction enzyme *EcoT22I* (manufactured by Takara Shuzo) to carry out the reaction at 30°C for 1 hour. The reaction solution was

fractionated by an agarose gel electrophoresis to recover about 2 µg of an *ApaI*-*EcoT22I* fragment of about 3.40 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

5 Next, 0.1 µg of the thus obtained *ApaI*-*EcoT22I* fragment of pBS_{hCyl} and 0.1 µg of the *ApaI*-*EcoT22I* of pBS_{ΔhCyl}-IL-2 were added to 10 µl in total volume of sterile water and ligated using DNA Ligation Kit Ver. 2 (manufactured by Takara Shuzo) in accordance with the
10 manufacture's instructions. Using the thus obtained recombinant plasmid DNA solution, the *E. coli* DH5α was transformed to obtain the plasmid pBS_{hCyl}-IL-2 shown in Fig. 22. The thus obtained plasmid (10 µg) was allowed to react in accordance with the manufacture's instructions
15 attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to an electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine nucleotide sequence of the inserted cDNA, and as a result, it was confirmed that a plasmid having the
20 nucleotide sequence of interest was obtained.

2. Stable expression of KM8871-hIL-2 using animal cells

(1) Construction of stable expression vector for KM8871-hIL-2

25 A stable expression vector for KM8871-hIL-2 was constructed as follows using the vector pKANTEX641HLC_{DRLm}-69 obtained in the item 6(1) of Example 1 for stable expression of anti-GD3 CDR-grafted antibody KM8871 and the plasmid pBS_{hCyl}-IL-2 obtained in the item 1(2) of Example 2
30 having cDNA encoding the fusion protein of hCyl with hIL-2.

 The plasmid pKANTEX641HLC_{DRLm}-69 (3 µg) obtained in the item 6(1) of Example 1 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed
35 with 10 units of a restriction enzyme *ApaI* (manufactured by

Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol and added to 10 µl of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and the solution was further mixed with 10 units of a restriction enzyme *Bam*HI (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an *Apa*I-*Bam*HI fragment of about 12.57 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 3 µg of each of the plasmid pBShCyl-IL-2 obtained in the item 1(2) of Example 2 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed with 10 units of a restriction enzyme *Apa*I (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol and added to 10 µl of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and the solution was further mixed with 10 units of a restriction enzyme *Bam*HI (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an *Apa*I-*Bam*HI fragment of about 1.45 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 0.1 µg of the thus obtained *Apa*I-*Bam*HI fragment of the plasmid pKANTEX641HLCDRIm-69 and 0.1 µg of the *Apa*I-*Bam*HI fragment of the plasmid pBShCyl-IL-2 were added to 10 µl in total volume of sterile water and ligated using DNA Ligation Kit Ver. 2 (manufactured by Pharmacia) in accordance with the manufacture's instructions. Using

the thus obtained recombinant plasmid DNA solution, an *E. coli* DH5 α was transformed to obtain the KM8871-hIL-2 stable expression vector pKANTEX8871-hIL2 shown in Fig. 23. The thus obtained plasmid (10 μ g) was allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to an electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence, and as a result, it was confirmed that a plasmid into which the DNA of interest was cloned was obtained.

(2) Expression of KM8871-hIL-2 in animal cells

Using 4 μ g of the KM8871-hIL-2 stable expression vector pKANTEX8871-hIL-2 obtained in the item 2(1) of Example 2, YB2/0 cell (ATCC CRL 1581) was transformed in accordance with the method described in the item 6(2) of Example 1, and selection was carried out finally with G418 (0.5 mg/ml) and MTX (200 nM) to obtain a transformed cell clone KM8871hIL2 showing an expression level of about 4 μ g/10⁶ cells/24 hours. Also, the KM8871hIL2 has been deposited on July 22, 1999, as FERM BP-6791 in National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology (Higashi 1-1-3, Tsukuba, Ibaraki, Japan).

(3) Purification of KM8871-hIL-2 from culture supernatant

In accordance with the method described in the item 6(3) of Example 1, the transformed cell clone KM8871hIL2 obtained in the item 2(2) of Example 2 capable of expressing KM8871-hIL-2 was cultured to obtain about 10.0 mg of purified KM8871-hIL-2 from about 3 L of the culture supernatant. A result of SDS-PAGE of the purified KM8871-hIL-2 is shown in Fig. 24. As shown in Fig. 24, molecular weight of the purified KM8871-hIL-2 was about 180 Kd under

non-reducing conditions, and two bands of about 65 Kd and about 25 Kd were found under reducing conditions. These molecular weights almost coincided with the molecular weights deduced from the cDNA nucleotide sequences of H chain of KM8871-hIL-2 and hIL-2 and L chain (H chain and hIL-2: about 64 Kd, L chain: about 24 Kd, whole molecule: about 176 Kd), and it was confirmed that the structure as antibody molecule was maintained after fusion of hIL-2.

3. Evaluation of activity of KM8871-hIL-2

(1) Reactivity of KM8871-hIL-2 with GD3 (ELISA method)

Reactivity of the purified KM8871-hIL-2 with GD3 (manufactured by DIA-IATRON) was measured in accordance with the method described in the item 2 of Example 1. In this case, a peroxidase-labeled goat anti-human IgG (H & L) antibody (manufactured by American Qualex, used by diluting 3,000 times with 1% BSA-PBS) was used as the secondary antibody solution. Fig. 25 shows a result of the examination of the reactivity carried out by fixing the amount of GD3 to be adsorbed to each well of a plate for ELISA use to 20 pmol/well and changing the concentration of the anti-GD3 CDR-grafted antibody KM8871 and KM8871-hIL-2 to be added. As shown in Fig. 25, it was found that KM8871-hIL-2 has a GD3 binding activity similar to or higher than that of the anti-GD3 CDR-grafted antibody KM8871. Fig. 26 shows a result of the examination of the reactivity of a constant concentration (10 µg/ml) of anti-GD3 CDR-grafted antibody KM8871 and KM8871-hIL-2, carried out by changing the kinds of ganglioside to be adsorbed to each well of a plate for ELISA use (adsorption amount: 20 pmol/well). As shown in Fig. 26, it was found that the KM8871-hIL-2 strongly binds to GD3 similar to the case of the anti-GD3 CDR-grafted antibody KM8871, and it was observed that KM8871-hIL-2 also strongly binds to GD2. As the cause of the cross-reactivity of KM8871-hIL-2 with GD2,

a possibility of the binding of hIL-2 with GD2, as well as a possibility of the change in binding specificity caused by a change in the three-dimensional structure of antibody V region due to fusion of hIL2, a possibility of the change
5 in the reactivity of secondary antibody caused by a change in the three-dimensional structure of antibody C region due to fusion of hIL2, and the like can be considered.

(2) Reactivity of KM8871-hIL-2 with cell surface GD3 (an
10 immunofluorescent method)

Reactivity of the purified KM8871-hIL-2 with human cancer cells was measured in accordance with the method described in the item 7(2) of Example 1. Results are shown in Fig. 27. As shown in Fig. 27, KM8871-hIL-2 showed a
15 strong reactivity with the human melanoma culture cell line G361, which was almost identical to that of the anti-GD3 CDR-grafted antibody KM8871. This result shows that the KM8871-hIL-2 is useful in the treatment and the like of GD3-positive human tumors including melanoma.

(3) Evaluation of hIL-2 activity of KM8871-hIL-2
20

Activity of the purified KM8871-hIL-2 as hIL-2 was measured in accordance with the following method. A mouse T cell line CTLL-2 (ATCC TIB214) of which growth shows
25 dependency on hIL-2 concentration was suspended in RPMI1640-FBS(10) medium at a density of 2×10^5 cells/ml and dispensed in 50 μ l/well into a 96 well microtiter plate (manufactured by Sumitomo Bakelite). A solution (50 μ l) prepared by diluting hIL-2 (manufactured by R & D SYSTEMS)
30 or purified KM8871-hIL-2 to various concentrations with RPMI1640-FBS(10) medium was added to each well and cultured at 37°C for 30 hours in a 5% CO₂ incubator. After the culturing the number of intact cells was counted using Cell Counting Kit (manufactured by Dojindo Laboratories) in
35 accordance with the manufacture's instructions. Results

are shown in Fig. 28. As shown in Fig. 28, KM8871-hIL-2 showed a biological activity to support CTLL-2 growth similar to that of hIL-2. These results show that the activity of KM8871-hIL-2 as hIL-2 is maintained after its fusion with the anti-GD3 CDR-grafted antibody KM8871.

(4) Activation of human effector cell and enhancement of cytotoxic activity by KM8871-hIL-2

In order to evaluate activation of human effector cells and accompanying enhancement of cytotoxic activity by the hIL-2 moiety of KM8871-hIL-2 *in vitro*, the cytotoxic activity was measured by the following method.

a. Preparation of target cell suspension

In accordance with the method described in "a" of the item 7(4) of Example 1, a human melanoma culture cell line G-361 was adjusted to a density of 2×10^5 cells/ml and used as the target cell suspension.

b. Preparation of human effector cell suspension

In accordance with the method described in "b" of the item 7(4) of Example 1, mononuclear cells were separated from venous blood of healthy person and re-suspended to a density of 5×10^5 cells/ml to be used as the effector cell suspension.

c. Activation of human effector cells

The effector cell suspension (100 μ l) prepared in "b" was dispensed into each well of a 96 well U bottom plate (manufactured by Falcon) (5×10^4 cells/well). Furthermore, 50 μ l of the anti-GD3 CDR-grafted antibody KM8871 or KM8871-hIL-2 was added thereto to give a final concentration of 11.1 nM and allowed to react at 37°C for 24 hours in a 5% CO₂ incubator.

d. Measurement of cytotoxic activity

The target cell suspension (50 μ l) prepared in "a" was added to each well of the plate prepared in "c" (1×10^4 cells/well). In this case, the ratio of the effector cells to the target cells becomes 5:1. After the reaction at 37°C for 4 hours, the plate was centrifuged and the amount of ^{51}Cr in the supernatant was measured by a γ -counter. The amount of the spontaneously dissociated ^{51}Cr was calculated by carrying out the same procedure using the medium alone instead of the effector cell suspension and antibody solution and measuring the amount of ^{51}Cr in the supernatant. The amount of the total dissociated ^{51}Cr was calculated by carrying out the same procedure by adding the medium alone instead of the antibody solution, and 1 N hydrochloric acid instead of the effector cell suspension, and measuring the amount of ^{51}Cr in the supernatant. The cytotoxic activity was calculated by the following equation.

$$\text{ADCC activity (\%)} = \frac{(\text{amount of } ^{51}\text{Cr in sample supernatant}) - (\text{amount of spontaneously released } ^{51}\text{Cr})}{(\text{amount of total } ^{51}\text{Cr}) - (\text{amount of spontaneously released } ^{51}\text{Cr})} \times 100$$

Results are shown in Fig. 29. As shown in Fig. 29, it was found that the cytotoxic activity induced by KM8871-hIL-2 was stronger than that induced by the anti-GD3 CDR-grafted antibody KM8871. This result shows that human effector cells can be activated and the cytotoxic activity can be enhanced by using KM8871-hIL-2. Thus, it was indicated that similar effects can be expected in its clinical application to human.

Example 3

Production of fusion protein of anti-GD3 chimeric antibody with human cytokine:

As an example of fusion protein of anti-GD3 chimeric antibody with human cytokine, a fusion protein of the anti-GD3 chimeric antibody KM871 with human IL-2, KM871-hIL-2, was prepared as follows to carry out evaluation of the activity.

1. Stable expression of KM871-hIL-2 using animal cells

(1) Construction of stable expression vector for KM871-hIL-2

A stable expression vector for KM871-hIL-2 was constructed as follows using the vector pKANTEX641 for stable expression of anti-GD3 chimeric antibody KM871 (Japanese Published Unexamined Patent Application No. 304989/93) and the plasmid pBShCyl-IL-2 obtained in the item 1(2) of Example 2 having cDNA encoding the fusion protein of hCyl with hIL-2.

The plasmid pKANTEX641 (3 µg) obtained in the item 6(1) of Example 1 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and 10 units of a restriction enzyme *Apa*I (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol and added to 10 µl of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and 10 units of a restriction enzyme *Bam*HI (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 30°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 2 µg of an *Apa*I-*Bam*HI fragment of about 12.57 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 3 µg of each of the plasmid pBShCyl-IL-2 obtained in the item 1(2) of Example 2 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol and added to 10 µl of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and 10 units of a restriction enzyme *BamHI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 2 µg of an *ApaI*-*BamHI* fragment of about 1.45 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 0.1 µg of the thus obtained *ApaI*-*BamHI* fragment of the plasmid pKANTEX641 and 0.1 µg of the *ApaI*-*BamHI* fragment of the plasmid pBShCyl-IL-2 were added to 10 µl in total volume of sterile water and ligated using DNA Ligation Kit Ver. 2 (manufactured by Pharmacia) in accordance with the manufacture's instructions. Using the thus obtained recombinant plasmid DNA solution, an *E. coli* DH5α was transformed to obtain the KM871-hIL-2 stable expression vector pKANTEX871-hIL2 shown in Fig. 30. The thus obtained plasmid (10 µg) was allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence, and as a result, it was confirmed that a plasmid into which the DNA of interest was cloned was obtained.

(2) Expression of KM871-hIL-2 in animal cells

Using 4 μ g of the KM871-hIL-2 stable expression vector pKANTEX871-hIL-2 obtained in the item 1(1) of Example 3, YB2/0 cell (ATCC CRL 1581) was transformed in accordance with the method described in the item 6(2) of Example 1, and selection was carried out finally with G418 (0.5 mg/ml) and MTX (200 nM) to obtain a transformed cell clone KM871hIL2 showing an expression level of about 4 μ g/ 10^6 cells/24 hours. Also, the KM871hIL2 has been deposited on October 19, 1999, as FERM BP-6918 in National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology (Higashi 1-1-3, Tsukuba, Ibaraki, Japan).

(3) Purification of KM871-hIL-2 from culture supernatant

In accordance with the method described in the item 6(3) of Example 1, the transformed cell clone KM871hIL2 obtained in the item 1(2) of Example 3 which expresses KM871-hIL-2 was cultured to obtain about 10.0 mg of purified KM871-hIL-2 from about 3 L of the culture supernatant. A result of SDS-PAGE of the purified KM871-hIL-2 is shown in Fig. 31. As shown in Fig. 31, molecular weight of the purified KM871-hIL-2 was about 180 Kd under non-reducing conditions, and two bands of about 65 Kd and about 25 Kd were found under reducing conditions. These molecular weights almost coincided with the molecular weights deduced from the cDNA nucleotide sequences of H chain of KM871-hIL-2 and hIL-2 and L chain (H chain and hIL-2: about 64 Kd, L chain: about 24 Kd, whole molecule: about 176 Kd), and it was confirmed that the structure as antibody molecule was maintained after fusion of hIL-2.

2. *In vitro* evaluation of KM871-hIL-2

(1) Reactivity of KM871-hIL-2 with GD3 (ELISA method)

Reactivity of the purified KM871-hIL-2 with GD3 (manufactured by DIA-IATRON) was measured in accordance with the method described in the item 2 of Example 1. In this case, a peroxidase-labeled goat anti-human IgG (H & L) antibody (manufactured by American Qualex, used by diluting 3,000 times with 1% BSA-PBS) was used as the secondary antibody solution. Fig. 32 shows results of the examination of the reactivity carried out by fixing the amount of GD3 to be adsorbed to each well of a plate for ELISA to 20 pmol/well and changing the concentration of the anti-GD3 chimeric antibody KM871 and KM871-hIL-2 to be added. As shown in Fig. 32, it was found that KM871-hIL-2 has a GD3 binding activity similar to or higher than that of the anti-GD3 chimeric antibody KM871. Fig. 33 shows results of the examination of the reactivity of a constant concentration (6.7 nM) of anti-GD3 chimeric antibody KM871 and KM871-hIL-2, carried out by changing the kinds of ganglioside to be adsorbed to each well of a plate for ELISA use (adsorption amount: 20 pmol/well). As shown in Fig. 33, it was found that the KM871-hIL-2 strongly binds to GD3 similar to the anti-GD3 chimeric antibody KM871, and it was confirmed that the addition of hIL-2 does not have large influence on the antigen specificity of KM871.

(2) Reactivity of KM871-hIL-2 with cell surface GD3 (immunofluorescent method)

Reactivity of the purified KM871-hIL-2 with human cancer cells was measured in accordance with the method described in the item 7(2) of Example 1. Results are shown in Fig. 34. As shown in Fig. 34, KM871-hIL-2 showed a strong reactivity with the human melanoma culture cell line G361 and SK-MEL-28, which was almost identical to that of the anti-GD3 chimeric antibody KM871. The results show

that the KM871-hIL-2 is useful in the treatment and the like of GD3-positive human tumors including melanoma.

(3) Evaluation of hIL-2 activity of KM871-hIL-2

5 Activity of the purified KM871-hIL-2 as hIL-2 was measured in accordance with the item 3(2) of Example 2, and results are shown in Fig. 35. As shown in Fig. 35, KM871-hIL-2 showed a biological activity to support CTLL-2 growth similar to that of hIL-2. The results show that the
10 activity of KM871-hIL-2 as hIL-2 is maintained after its fusion with the anti-GD3 chimeric antibody KM871.

(4) Activation of human lymphocyte by KM871-hIL-2

15 In order to evaluate activation of human lymphocyte and accompanying enhancement of cytotoxic activity by the hIL-2 moiety of KM871-hIL-2 in vitro, the cytotoxic activity was measured by the following method in which the method described in the item 7(4) of Example 1 was partially changed.

20

a. Preparation of target cell suspension

 In accordance with the method described in "a" of the item 7(4) of Example 1, a human melanoma culture cell line G-361 was adjusted to give a density of 2×10^5 cells/ml
25 and used as the target cell suspension.

b. Preparation of effector cell suspension

 In accordance with the method described in "b" of the item 7(4) of Example 1, mononuclear cells were
30 separated from venous blood of healthy person and re-suspended to give a density of 1×10^7 cells/ml to be used as the effector cell suspension.

c Activation of lymphocyte

The effector cell suspension (50 μ l) prepared in "b" was dispensed into each well of a 96 well U bottom plate (manufactured by Falcon) (5×10^4 cells/well).
5 Furthermore, 50 μ l of RPMI1640-FBS(10) medium to which the anti-GD3 chimeric antibody KM871 or KM871-hIL-2 had been added was added thereto to give a final concentration of 1 nM (molecular weight of KM871 was calculated as 150 Kd, and molecular weight of KM871-hIL-2 as 180 Kd) and incubated at
10 37°C for 72 hours in a 5% CO₂ incubator.

d. Measurement of cytotoxic activity

The target cell suspension (50 μ l) prepared in "a" was added to each well of the plate prepared in
15 "c" (1×10^4 cells/well). In this case, the ratio of the effector cells to the target cells becomes 5:1. Furthermore, 50 μ l of RPMI1640-FBS(10) medium to which the anti-GD3 chimeric antibody KM871 or KM871-hIL-2 had been added was added thereto to give a final concentration of 1
20 nM to carry out the reaction at 37°C for 4 hours, and then the plate was centrifuged and the amount of ⁵¹Cr in the supernatant was measured by a γ -counter. The amount of the spontaneously dissociated ⁵¹Cr was calculated by carrying out the same procedure using the medium alone instead of
25 the effector cell suspension and antibody solution and measuring the amount of ⁵¹Cr in the supernatant. The amount of the total dissociated ⁵¹Cr was calculated by carrying out the same procedure by adding the medium alone instead of the antibody solution, and 1 N hydrochloric acid instead
30 of the effector cell suspension, and measuring the amount of ⁵¹Cr in the supernatant. The cytotoxic activity was calculated by the method described in "b" of the item 7(4) of Example 1, and results are shown in Fig. 36. As shown in Fig. 36, it was found that a cytotoxic activity induced
35 by KM871-hIL-2 was stronger than that induced by KM871.

Based on the results, it is expected that the KM871-hIL-2 will also enhance therapeutic effects of KM871 in the clinical application of GD3-positive human tumors by the activation of human lymphocyte via the hIL-2 moiety.

5

3. In vivo evaluation of KM871-hIL-2

(1) Production of B16 cell transfectants introduced with GD3 synthase gene for mouse syngenic model

10 a. Construction of GD3 synthase gene stable expression vector

A vector pAMo-GD3 (3 µg; WO 94/23020) for expressing GD3 synthase gene in animal cell was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 50 mM sodium chloride, 10 mM magnesium chloride and 1 mM DTT, and
15 10 units of a restriction enzyme *Hind*III (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol and added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride,
20 10 mM magnesium chloride, 1 mM DTT, 100 µg/ml BSA and 0.01% Triton X-100, and 10 units of a restriction enzyme *Not*I (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, and the 5'
25 protruding end formed by the restriction enzyme digestion was changed to blunt end using DNA Blunting Kit (manufactured by Takara Shuzo). The reaction solution was fractionated by agarose gel electrophoresis to recover about 2 µg of a blunt-ended *Hind*III-*Not*I fragment of about
30 2.1 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 3 µg of the anti-GD3 chimeric antibody expression vector pKANTEX641 was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride,
35 10 mM magnesium chloride and 1 mM DTT, and 10 units of a

restriction enzyme *EcoRI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, and the 5' protruding end formed by the restriction enzyme digestion was changed to blunt end using DNA Blunting Kit (manufactured by Takara Shuzo). The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5) and 10 mM magnesium chloride, and the solution was further mixed with 10 units of a 5'-terminal dephosphorylation enzyme, calf intestine-derived alkaline phosphatase (manufactured by Takara Shuzo), and allowed to react at 37°C for 1 hour for dephosphorylation of the 5' end. The reaction solution was subjected to phenol treatment and then fractionated by agarose gel electrophoresis to recover about 2 µg of a blunt end-dephosphorylated *EcoRI*-*NruI* fragment of about 9.4 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 0.1 µg of the *EcoRI*-*NruI* fragment derived from the plasmid pKANTEK641 and 0.1 µg of the *HindIII*-*NotI* fragment derived from the plasmid pAMoGD3, both obtained in the above, were added to 10 µl in total volume of sterile water and ligated using DNA Ligation Kit Ver. 2 (manufactured by Takara Shuzo) in accordance with the manufacture's instructions. Using the thus obtained recombinant plasmid DNA solution, an *E. coli* DH5α was transformed to obtain the GD3 synthase-stable expression vector pKANTEKGD3 shown in Fig. 37.

b. Introduction of GD3 synthase-expression vector into B16 cells

Using 4 µg of the GD3 synthase gene-stable expression vector pKANTEKGD3 obtained in "a" of the item 3(1) of Example 3, B16-F10 cell (ATCC CRL 1581) was transformed in accordance with the method described in the

item 6(2) of Example 1, and selection was carried out finally by G418 (0.5 mg/ml) to obtain transfectant clones B16 8-3 and B16 29-10.

5 c. Analysis of GD3 expression level of GD3-expressing line by immunofluorescent method

Each of 1×10^6 cells of the GD3 synthase gene transfectants B16 8-3 and B16 29-10 obtained in "b" of the item 3(1) of Example 3 and the parent line B16 F10 cells
10 were suspended in PBS, put into microtubes and centrifuged (2,000 rpm for 2 minutes), and the thus washed cells were stirred after adding 50 μ l of the anti-GD3 chimeric antibody KM871 (a solution prepared by adjusting to 5 μ g/ml with 1% BSA-PBS, negative control was 1% BSA-PBS alone) and
15 then allowed to react at 4°C for 1 hour. After the reaction, followed by centrifugation three times with PBS for washing, the cells were mixed with 100 μ l of biotin-labeled goat anti-human IgG (H - L) solution (manufactured by VECTOR, used by diluting 400 times with 1% BSA-PBS),
20 stirred and then allowed to react at 4°C for 1 hour. After centrifugation three times with PBS for washing, 100 μ l of a streptavidin-labeled Cy5 pigment solution (Streptavidin-RED670 manufactured by GIBCO, used by diluting 100 times with 1% BSA-PBS) was added and stirred and then allowed to
25 react at 4°C for 1 hour. After the reaction, followed by centrifugation three times with PBS for washing, the cells were again suspended in PBS to carry out the analysis using a flow cytometer EPICS Elite (manufactured by COULTER). Results are shown in Fig. 38. As shown in Fig. 38, the
30 anti-GD3 chimeric antibody KM871 reacted only with B16 8-3 and B16 29-10 but did not show reactivity with the parent line B16 F10 cells. The above results show that both of the transfectants B16 8-3 and B16 29-10 express GD3 on the surface and can be used as a model for measuring *in vivo*

antitumor effects of KM871-hIL-2 by transplanting them into C57 BL/6 mouse as the origin of B16 cells.

(2) Measurement of in vivo antitumor effects of KM871-hIL-2

- 5 a. Evaluation with mice transplanted with GD3-expressing transformant B16-29-10

10 The GD3-expressing B16 cell B16 29-10 obtained in "b" of the item 3(1) of Example 3 was suspended in PBS to give a density of 2.5×10^6 cells/ml, and 100 μ l of the suspension was injected into a C57 BL/6 Cr slc mouse (female, 8 week-old, available from Japan SLC) through its tail vein. Also, starting on the day of tumor transplantation, 50 or 200 μ g/day of KM871 or 20 or 50 μ g/day of KM871-hIL-2 was administered intravenously once a day continuously for 5 days. The lungs were excised on the 15th day after the tumor transplantation, and visible black metastatic foci on the surface were counted, and results are shown in Table 3 and Fig. 39.

20

Table 3

Group		The number of metastatic foci on the lung surface	Average value
Non-administered group		147/136/106/72/70	106.2
KM871-hIL-2	20 μ g	20/15/11/2/2/1	8.5
KM871-hIL-2	50 μ g	15/12/5/2/0/0	5.7
KM871	50 μ g	185/114/86/77	115.5
KM871	200 μ g	113/87/86/86/77/55	84

As shown in Table 3 and Fig. 39, KM871-hIL-2 showed clearly more potent anti-metastatic effect than that of KM871 in the model mouse of the clone B16-29-10.

25

- b. Evaluation with mice transplanted with GD3-expressing transformant B16-8-3

Using the GD3-expressing B16 cell B16-8-3 obtained in "b" of the item 3(1) of Example 3, anti-metastatic effect of KM871-hIL-2 was measured in accordance with the method described in "a" of the item 3(2) of Example 3. The dose was 40 or 100 µg/day for both KM871 and KM871-hIL-2. Results are shown in Table 4 and Fig. 40.

Table 4

Group	The number of metastatic foci on the lung surface	Average value
Non-administered group	170/128/113/66/42/16/0	76.4
KM871-hIL-2 40 µg	1/0/0	0.3
KM871-hIL-2 100 µg	3/0/0	1
KM871 40 µg	128/70/30	76
KM871 100 µg	26/20/17	21

As shown in Table 4 and Fig. 40, KM871-hIL-2 showed more potent anti-metastatic effect than that of KM871 in the model mouse of the clone B16-8-3.

c. Comparison with combined administration of KM871 and hIL-2

The GD3-expressing B16 cells B16-29-10 obtained in "b" of the item 3(1) of Example 3 were suspended in PBS to give a density of 2.5×10^6 cells/ml, and 100 µl of the suspension was injected into a C57 BL/6 Cr slc mouse (female, 8 week-old, available from Japan SLC) through its tail vein. Also, starting on the day of tumor transplantation, 6 µg/day (0.0333 nmol) or 18 µg/day (0.1 nmol) of KM871-hIL-2 was administered intravenously once a day continuously for 5 days. Using other groups, a mixed solution of KM871 and hIL-2 (manufactured by Peptotech) corresponding to the amount of the administered KM871-hIL-2 (5 µg (0.0333 nmol) of KM871 and 1 µg (0.0667 nmol) of hIL-2, or 15 µg (0.1 nmol) of KM871 and 3 µg (0.2 nmol) of hIL-

2) was administered through the tail vein once a day continuously for 5 days, starting on the day of tumor transplantation. The lungs were excised on the 15th day after the tumor transplantation, and visible black metastatic foci on the surface were counted, and results are shown in Table 5 and Fig. 41.

Table 5

Group		The number of metastatic foci on the lung surface	Average value
Non-administered group		163/132/97/95/88/57	105.3
KM871-hIL-2	6 μ g	26/26/17/12/8	17.8
KM871-hIL-2	18 μ g	43/34/28/13/2	24
KM871 + hIL-2	5 μ g + 1 μ g	215/117/97/85/56	114
KM871 + hIL-2	15 μ g + 3 μ g	92/61/48/46/42	57.8

As shown in Table 5 and Fig. 41, KM871-hIL-2 showed more potent anti-metastatic effect than that of the combined administration of the equivalent amount of KM871 and hIL-2 in the model mouse of the clone B16-29-10. The results show that the *in vivo* antitumor effect shown by KM871-hIL-2 is not a simple additive effect of KM871 and hIL-2 but a result of the efficient induction of antitumor immunity due to activation of peripheral lymphocytes by the hIL-2 moiety of KM871-hIL-2 accumulated in the tumor, thus showing a possibility that KM871-hIL-2 will become a therapeutic agent having higher therapeutic effect than that of the conventional administration of hIL-2 alone or its combination use with an antibody and also having reduced side effects derived from hIL-2.

d. Evaluation by solid tumor early model of B16-29-10 cell

The GD3-expressing B16 cell B16-29-10 obtained in "b" of the item 3(1) of Example 3 was suspended in PBS to

give a density of 1×10^7 cells/ml, and 50 μ l of the suspension was transplanted subcutaneously in C57 BL/6 mice (male, 7 week-old, available from Charles River Japan). By setting the following administration groups, each agent was administered intravenously once a day continuously for 5 days, starting on the day of tumor transplantation.

Non-administered group

- hIL-2: 10 μ g/day (0.667 nmol)
- 10 KM871: 50 μ g/day (0.333 nmol)
- KM871-hIL-2: 60 μ g/day (0.333 nmol)

The test was carried out using 5 animals for each group. Each agent was prepared to give a concentration of 200 μ l/animal by diluting with a citrate buffer. Five day after the transplantation, the tumor diameter was measured periodically using slide calipers, and the antitumor effect was judged based on the ratio of the average value of tumor volumes in each administered group to the average value of tumor volumes in non-administered group and on the number of survived days after commencement of the administration. The tumor volume was calculated by the following equation

$$\text{Tumor volume} = (\text{width})^2 \times \text{length} \times 0.5$$

25

The average value of tumor volumes in each group is shown in Table 6, results of the ratio of the average value of tumor volumes in each group to the average value of tumor volumes in non-administered group are shown in Table 7 and Fig. 42, and results of the number of survived days are shown in Table 8.

30

Table 6

Days after administration	Non-administered group	KM871	hIL-2	KM871-hIL-2
0	0	0	0	0
5	75	60	36	49
7	131	82	67	34
9	306	146	115	87
11	756	494	359	231
12	1325	939	489	332
14	2333	1701	1147	733
16	3656	3331	2268	1432
20	7799	8345	5765	3381

Unit is mm³

Table 7

Days after administration	Non-administered group	KM871	hIL-2	KM871-hIL-2
0	1.000	1.000	1.000	1.000
5	1.000	0.796	0.484	0.656
7	1.000	0.624	0.510	0.261
9	1.000	0.477	0.376	0.285
11	1.000	0.654	0.475	0.305
12	1.000	0.709	0.369	0.251
14	1.000	0.729	0.492	0.314
16	1.000	0.911	0.620	0.392
20	1.000	1.070	0.739	0.434

Table 8		
Group	Survived days after commencement of administration (day)	Average value (day)
Non-administered group	24/27/27/30/30	27.6
hIL-2	26/28/32/32/33	30.2
KM871	27/30/30/32/35	30.8
KM871-hIL-2	32/32/43/51/57	43.0

As shown in Tables 6, 7 and 8 and Fig. 42, KM871-hIL-2 showed more potent growth inhibition effect and life-prolonging effect than those of the hIL-2 alone or antibody alone.

e. Evaluation by solid tumor advanced stage model of B16-29-10 cell

The GD3-expressing B16 cell B16-29-10 was suspended in PBS to a density of 1×10^8 cells/ml, and 50 μ l of the suspension was transplanted subcutaneously in C57 BL/6 mice (male, 7 week-old, available from Charles River Japan). By setting the following administration groups, tumor volumes on the 6th day after the tumor transplantation were calculated by the measuring method described in "d" of the item 3(2) of Example 3 to select individuals within the range of from 10 to 80 mm³, and then each of the following agents was administered intravenously once a day continuously for 8 days.

Non-administered group

hIL-2: 10 μ g/day (0.667 nmol)

KM871: 50 μ g/day (0.333 nmol)

KM871: 200 μ g/day (1.33 nmol)

KM871-hIL-2: 24 μ g/day (0.133 nmol)

KM871-hIL-2: 60 μ g/day (0.333 nmol)

The test was carried out using 3 animals only in the KM871-hIL-2 60 µg/day administration group, and 5 animals for each of the other groups. Each agent was prepared to a concentration of 200 µl/animal by diluting with citrate buffer. Starting on the day of transplantation, the tumor diameter was measured periodically using slide calipers, and the antitumor effect was judged based on the ratio of the average value of tumor volumes on the measured day and tumor volumes on the day of commencement of the administration in each mouse in each treated groups (hereinafter referred to as "V/V0") to the V/V0 in non-administered group and on the number of survived days after commencement of the administration.

Results of V/V0 in each group are shown in Table 9, results of the ratio of the V/V0 in each group to the V/V0 in non-administered group are shown in Table 10 and Fig. 43, and results of the number of survived days are shown in Table 11.

Table 9

Days	Non-administered group	hIL-2 10 µg	KM871 50 µg	KM871 200 µg	KM871-hIL-2 24 µg	KM871-hIL-2 60 µg
0	1.00	1.00	1.00	1.00	1.00	1.00
2	2.15	3.03	2.57	1.86	2.09	1.54
4	5.36	6.15	6.37	4.98	3.79	2.88
6	13.18	14.31	14.43	9.35	8.12	6.85
7	16.45	17.68	18.99	11.22	10.05	8.58
8	21.09	21.31	24.05	16.56	13.04	11.50
9	28.95	28.42	33.43	21.08	16.10	15.19
11	54.97	48.52	56.07	42.65	29.13	27.01
13	93.70	86.85	100.34	74.16	58.17	49.80

Table 10

Days	Non-administered group	hIL-2 10 µg	KM871 50 µg	KM871 200 µg	KM871-hIL-2 24 µg	KM871-hIL-2 60 µg
0	1.000	1.000	1.000	1.000	1.000	1.000
2	1.000	1.406	1.196	0.862	0.973	0.718
4	1.000	1.147	1.187	0.928	0.706	0.537
6	1.000	1.086	1.096	0.710	0.616	0.520
7	1.000	1.075	1.155	0.682	0.611	0.522
8	1.000	1.011	1.141	0.785	0.619	0.545
9	1.000	0.982	1.155	0.728	0.556	0.525
11	1.000	0.883	1.020	0.776	0.530	0.491
13	1.000	0.927	1.071	0.791	0.621	0.531

Table 11

Group	Survived days after commencement of administration (day)	Average value (day)
Non-administered group	15/23/23/24/34	23.8
hIL-2	15/15/18/27/29	20.8
KM871 50 µg	19/20/23/30/34	25.2
KM871 200 µg	18/19/23/27/34	24.2
KM871-hIL-2 24 µg	29/38/39/41/49	39.2
KM871-hIL-2 60 µg	29/32/37	32.7

5 As shown in Tables 9, 10 and 11 and Fig. 43, KM871-hIL-2 showed more potent growth inhibition effect and life-prolonging effect than those of the hIL-2 alone or antibody alone.

10 f. Effect of combination use of KM871 and KM871-hIL-2 in solid tumor advanced stage model of B16-29-10 cell

15 The GD3-expressing B16 cell B16-29-10 obtained in "b" of the item 3(1) of Example 3 was suspended in PBS to give a density of 1×10^8 cells/ml, and 50 µl of the suspension was transplanted subcutaneously in C57 BL/6 mice

(male, 7 week-old, available from Charles River Japan). By setting the following administration groups, tumor volumes on the 6th day after the tumor transplantation (0th day) were calculated by the measuring method described in d. of the item 3(2) of Example 3 to select individuals within the range of from 10 to 100 mm³, and then each of the following agents was administered intravenously.

Non-administered group

- 10 hIL-2: 10 µg/day × 5 days continuous
administration (0th to 4th days)
- KM871: 800 µg/day × single administration
(0th day)
- 15 ·KM871 + KM871-hIL-2: 800 µg/day × single administration
(0th day) + 15 µg/day × 5 days
continuous administration (0th to
4th days)
- 20 ·KM871 + KM871-hIL-2: 800 µg/day × single administration
(0th day) + 60 µg/day × 5 days
continuous administration (0th to
4th days)

The test was carried out using 5 animals for each group. Each agent was prepared to give a concentration of 200 µl/animal by diluting with a citrate buffer. Starting on the day of transplantation, the tumor diameter was measured periodically using slide calipers, and the antitumor effect was judged based on the ratio of the V/V₀ in each treated group to the V/V₀ in non-administered group and on the number of survived days after commencement of the administration. Results of V/V₀ in each group are shown in Table 12, results of the ratio of the V/V₀ in each group to the V/V₀ in non-administered group are shown in Table 13 and Fig. 44, and results of the number of survived days are shown in Table 14.

Table 12

Days	Non-administered group	hIL-2 10 µg × 5	KM871 800 µg × 1	KM871 + KM871-hIL-2 800 µg × 1 + 15 µg × 5	KM871 + KM871-hIL-2 800 µg × 1 + 60 µg × 5
0	1.00	1.00	1.00	1.00	1.00
2	3.09	2.76	1.81	1.68	1.49
4	8.54	4.10	4.10	2.56	2.68
6	17.48	9.96	14.17	6.42	6.03
8	46.84	20.63	25.08	11.62	10.58
10	63.99	33.09	49.33	22.98	17.65
12	104.89	57.06	75.47	38.46	27.76

Table 13

Days	Non-administered group	hIL-2 10 µg × 5	KM871 800 µg × 1	KM871 + KM871-hIL-2 800 µg × 1 + 15 µg × 5	KM871 + KM871-hIL-2 800 µg × 1 + 60 µg × 5
0	1.00	1.00	1.00	1.00	1.00
2	1.00	0.89	0.59	0.54	0.48
4	1.00	0.48	0.48	0.30	0.31
6	1.00	0.57	0.81	0.37	0.34
8	1.00	0.44	0.54	0.25	0.23
10	1.00	0.52	0.77	0.36	0.28
12	1.00	0.54	0.72	0.37	0.26

5

Table 14

Group	Survived days after commencement of administration (day)	Average value (day)
Non-administered group	9/12/16/16/29	16.4
hIL-2	17/17/24/32/36	25.2
KM871	8/16/17/26/31	19.6
KM871 + KM871-hIL-2 15 µg	17/24/29/32/43	29.7
KM871 + KM871-hIL-2 60 µg	29/29/36/36/43	34.6

As shown in Table 12, Table 13, Fig. 44 and Table 14, combined administration of KM871 and KM871-hIL-2 showed more potent growth inhibition effect and life-prolonging effect than those of the hIL-2 alone or antibody alone. The results show a possibility that combined administration of KM871-hIL-2 and KM871 will become an excellent therapeutic method for GD3-positive cancers.

4. Analysis of *in vivo* antitumor mechanism of KM871-hIL-2

(1) Activation of mouse effector cell and enhancement of cytotoxic activity by KM871-hIL-2

In order to evaluate activation of mouse effector cell by the hIL-2 moiety of KM871-hIL-2 and accompanying enhancement of cytotoxic activity, the cytotoxic activity was measured in accordance with the method shown below.

a. Preparation of mouse effector cell suspension

The following administration groups were set using C57 BL/6 mice (female, 8 week-old, Japan SLC), and each agent was administered intravenously.

Non-administered group

hIL-2: 10 μ g (0.667 nmol)/day \times 5 days of continuous administration (0th to 4th days)

KM871-hIL-2: 60 μ g (0.333 nmol)/day \times single administration (2nd day)

KM871-hIL-2: 60 μ g (0.333 nmol)/day \times two administrations (0th day and 2nd day)

After completion of all administrations on the 4th day, the spleen was excised, mashed using a slide glass in 6 ml of RPMI1640-FBS(10) medium and then passed through a 70 μ m diameter nylon mesh (manufactured by Falcon) to prepare a suspension of splenocytes. A density separation medium for mouse lymphocytes (5 ml), Lympholite-M, was put

into a 15 ml tube, overlaid gently with 5 ml of the thus obtained splenocyte suspension and then centrifuged at room temperature for 20 minutes at $1,000 \times g$. The lymphocyte layer of the interface was recovered using a dropping pipette, 10 ml of RPMI1640-FBS(10) medium was added thereto, and the mixture was centrifuged ($800 \times g$, 4°C , 10 minutes) and then the supernatant was discarded for washing. After repeating the washing procedure again, an effector cell suspension was prepared to give a final density of 2×10^7 cells/ml by adding RPMI1640-FBS(10) medium.

b. Preparation of target cell suspension

In accordance with the method described in "a" of the item 7(4) of Example 1, the GD3-expressing B16 cell B16-29-10 obtained in "b" of the item 3(1) of Example 3 was labeled with ^{51}Cr to give 2×10^5 cells/ml of a target cell suspension.

c. Measurement of cytotoxic activity

Each of 100 μl of the effector cell suspension prepared in "a", 50 μl of the target cell suspension prepared in "b" (1×10^4 cells/well) and KM871 serially diluted with RPMI1640-FBS(10) medium to give a final concentration of 0, 0.05, 0.5, 5 or 50 $\mu\text{g/ml}$ was added to each well of a 96 well U bottom plate. In this case, the ratio of the effector cells to the target cells becomes 200:1. After the reaction at 37°C for 10 hours, the plate was centrifuged and the amount of ^{51}Cr in the supernatant was measured by a γ -counter. The amount of the spontaneously dissociated ^{51}Cr was calculated by carrying out the same procedure using the medium alone instead of the effector cell suspension and antibody solution and measuring the amount of ^{51}Cr in the supernatant. The amount of the total dissociated ^{51}Cr was calculated by carrying out the same procedure by adding the medium alone instead

of the antibody solution, and 1 N hydrochloric acid instead of the effector cell suspension, and measuring the amount of ^{51}Cr in the supernatant. The cytotoxic activity was calculated by the method described in "b" of the item 7(4) of Example 1. Results are shown in Fig. 45. As shown in Fig. 45, it was found that KM871-hIL-2 induces the cytotoxic activity higher than the 5 times administration of hIL-2, by both of its single and two times administrations.

(2) Analysis of effector cells activated by KM871-hIL-2 (immunofluorescent method)

In order to measure influences of the administration of KM871-hIL-2 on effector cells relating to the antitumor effect in the mouse model, a flow cytometer analysis was carried out by the following procedure. The GD3-expressing B16 cell B16-29-10 obtained in b. of the item 3(1) of Example 3 was suspended in PBS to a density of 1×10^7 cells/ml, and 50 μl of the suspension was transplanted under the ventral skin of a C57 BL/6 mouse (male, 7 week-old, available from Charles River Japan). By setting the following administration groups, each agent was administered intravenously once a day continuously for 5 days, starting on the day of tumor transplantation.

Non-administered group

hIL-2: 10 $\mu\text{g/day}$ (0.667 nmol)

KM871-hIL-2: 60 $\mu\text{g/day}$ (0.333 nmol)

Regarding KM871-hIL-2 (60 $\mu\text{g/day}$), a group in which the frequency of administration was reduced to two times (on the day of transplantation and the 3rd day after transplantation) was also set. The test was carried out using two animals for each group. After completion of all administrations on the 4th day after transplantation, the

spleen was excised, mashed using a slide glass in 6 ml of RPMI1640-FBS(10) medium and then passed through a 70 μ m diameter nylon mesh (manufactured by Falcon) to prepare a suspension of splenocytes. A density separation medium for mouse lymphocytes (5 ml), Lympholite-M, was put into a 15 ml tube, overlaid gently with 5 ml of the thus obtained splenocyte suspension and then centrifuged at room temperature for 20 minutes at 1,000 \times g. The lymphocyte layer of the interface was recovered using a dropping pipette, 10 ml of RPMI1640-FBS(10) medium was added thereto, and the mixture was centrifuged (800 \times g, 4°C, 10 minutes) and then the supernatant was discarded for washing. After repeating the washing procedure again, a spleen lymphocyte suspension was prepared to give a final density of 3×10^7 cells/ml by adding 1% BSA-PBS containing 5% mouse serum (manufactured by Chemicon). The lymphocyte suspension was dispensed in 50 μ l into a 96 well U bottom plate and each of the following fluorescence-labeled antibodies was further added thereto to carry out the reaction at 4°C for 1 hour.

Phycoerythrin (hereinafter referred to as "PE")-labeled rat anti-mouse CD4 antibody (manufactured by CEDERLANE):	3 μ l
FITC-labeled rat anti-mouse CD8 antibody (manufactured by Serotec):	5 μ l
FITC-labeled rat anti-mouse NK cell antibody (manufactured by Serotec):	15 μ l
FITC-labeled rat anti-mouse macrophage antibody (manufactured by Serotec):	10 μ l
PE-labeled rat anti-mouse neutrophil antibody (manufactured by Caltac):	5 μ l

By analyzing using a flow cytometer EPICS ELITE (manufactured by Coulter), the ratio of a group stained with each antibody to the total intact cells was calculated,

and results are shown in Table 15 and Fig. 46. Each numeral in the table shows the average value of two animals in each group (unit: %).

5

Table 15

	CD4+T cell	CD8+T cell	NK cell	Macrophage	Neutrophil
No treatment	28.5	21.7	3.71	0.643	7.63
hIL-2 5 x administration	21.3	20.7	16.2	0.448	13.9
KM871-hIL-2 5 x administration	16.1	20.9	18.5	0.392	7.26
KM871-hIL-2 2 x administration	27.2	28.8	17.0	0.498	6.06

As shown in Table 15 and Fig. 46, increase in the ratio of NK cells in spleen lymphocytes was observed by the administration of both hIL-2 and KM871-hIL-2. The results show a possibility that the antitumor effect of KM871-hIL-2 is enhanced via the proliferation of NK cells.

(3) Effector cell analysis of antitumor effect using in vivo depletion method (in vitro cytotoxic activity)

Mechanism analysis of the antitumor effect of KM871-hIL-2 was carried out by the method shown below, using mice prepared by administering antibodies against NK cell, CD4-positive T cell (helper T cell) and CD8-positive T cell (cytotoxic T cell), as IL-2 responsive effector cells in antitumor immunity, into mice to thereby deplete respective cell group from the animal body. The GD3-expressing B16 cell B16-29-10 obtained in "b" of the item 3(1) of Example 3 was suspended in PBS to a density of 5×10^6 cells/ml, and 100 μ l of the suspension was injected intravenously in each of C57 BL/6 mice (male, 7 week-old, available from Charles River Japan). Also, KM871-hIL-2 was administered intravenously at a dose of 24 μ g/day once a

day for continuous 5 days (0th day to 4th day) starting on the day of tumor transplantation (0th day). At the same time, no administration group was also set as a control group. In addition, KM871-hIL-2 administration groups and no administration groups were also set in the same manner on mice into which each of anti-mouse NK1.1 antibody, anti-mouse CD4 antibody and anti-mouse CD8 antibody was administered by intraperitoneal injection at a dose of 0.5 mg/day three times (-3rd day, 1st day and 3rd day). Five days after the tumor transplantation, the spleen was excised and the cytotoxic activity was measured by the method described in the item 4(1) of Example 3. In this case, the concentration of KM871 at the time of assay was fixed to 10 µg/ml, and the assay time was set to 4 hours. Results are shown in Fig. 47. As shown in Fig. 47, it was found that the main effector cell of the antitumor effect of KM871-hIL-2 in the mice model is NK cell.

(4) Effector cell analysis of antitumor effect using *in vivo* depletion method (*in vivo* antitumor activity)

Mechanism analysis of the antitumor effect of KM871-hIL-2 was carried out by the method shown below, using mice prepared by administering antibodies against NK cell, CD4-positive T cell (helper T cell) and CD8-positive T cell (cytotoxic T cell), as IL-2 responsive effector cells in antitumor immunity, into mice to thereby deplete respective cell group from the animal body. The GD3-expressing B16 cell B16-29-10 obtained in "b" of the item 3(1) of Example 3 was suspended in PBS to give a density of 5×10^6 cells/ml, and 100 µl of the suspension was injected intravenously in each of C57 BL/6 mice (male, 7 week-old, manufactured by Charles River Japan). Also, KM871-hIL-2 was administered intravenously at a dose of 24 µg/day once a day for continuous 5 days (0th day to 4th day) starting on the day of tumor transplantation (0th day). At the same

time, no administration group was also set as a control group. In addition, KM871-hIL-2 administration groups and no administration groups were also set in the same manner on mice into which each of anti-mouse NK1.1 antibody, anti-mouse CD4 antibody and anti-mouse CD8 antibody were administered by intraperitoneal injection at a dose of 0.5 mg/day four times (-3rd day, 1st day, 3rd day and 7th). On the 15th day after tumor transplantation, the lungs were excised and the visible black metastatic foci on the surface were counted, and results are shown in Table 16 and Fig. 48.

Table 16

Admin- istra- tion	Elimi- nation	Number of metastatic foci							Average value	Metastasis inhibition ratio (%)
None	No	302	274	269	234	177	176	173	229.3	0.0
	NK cell	443	430	389	380	373	361	359	390.7	0.0
	CD4+T cell	291	271	219	211	197	176	115	198.9	0.0
	CD8+T cell	266	226	191	191	182	144	73	181.9	0.0
KM871- hIL-2 24 µg	No	77	54	46	46	36			51.8	77.4
	NK cell	422	384	358	356	352			374.4	4.2
	CD4+T cell	164	134	103	95	47			108.6	45.4
	CD8+T cell	88	67	53	35	21			52.8	71.0

As shown in Table 16 and Fig. 48, it was found that the main effector cell of the antitumor effect of KM871-hIL-2 in the model animals is NK cell. Also, since partial inhibition of the antitumor effect was found by the elimination of CD4-positive T cell, a possibility was suggested that CD4-positive T cell is indirectly concerned in the antitumor effect of KM871-hIL-2 by activating NK cell.

INDUSTRIAL APPLICABILITY

According to the present invention, a human CDR-grafted antibody which specifically reacts with GD3 or the antibody fragment thereof and derivatives of an antibody
5 against ganglioside GD3 or the antibody fragment thereof are provided.

Free Text of Sequence Listings

10 SEQ ID NO:9-Explanation of artificial sequence: Amino acid sequence obtained by synthetic DNA
SEQ ID NO:10-Explanation of artificial sequence: Amino acid sequence obtained by synthetic DNA
SEQ ID NO:11-Explanation of artificial sequence: Synthetic DNA
15 SEQ ID NO:12-Explanation of artificial sequence: Synthetic DNA
SEQ ID NO:13-Explanation of artificial sequence: Synthetic DNA
SEQ ID NO:14-Explanation of artificial sequence: Synthetic
20 DNA
SEQ ID NO:15-Explanation of artificial sequence: Synthetic DNA
SEQ ID NO:16-Explanation of artificial sequence: Synthetic DNA
25 SEQ ID NO:17-Explanation of artificial sequence: Synthetic DNA
SEQ ID NO:18-Explanation of artificial sequence: Synthetic DNA
SEQ ID NO:19-Explanation of artificial sequence: Synthetic
30 DNA
SEQ ID NO:20-Explanation of artificial sequence: Synthetic DNA
SEQ ID NO:21-Explanation of artificial sequence: Synthetic DNA

SEQ ID NO:22-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:23-Explanation of artificial sequence: Synthetic
DNA
5 SEQ ID NO:24-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:25-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:26-Explanation of artificial sequence: Synthetic
10 DNA
SEQ ID NO:27-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:28-Explanation of artificial sequence: Synthetic
DNA
15 SEQ ID NO:29-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:30-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:31-Explanation of artificial sequence: Synthetic
20 DNA
SEQ ID NO:32-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:33-Explanation of artificial sequence: Synthetic
DNA
25 SEQ ID NO:34-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:35-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:36-Explanation of artificial sequence: Synthetic
30 DNA
SEQ ID NO:37-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:38-Explanation of artificial sequence: Synthetic
DNA

SEQ ID NO:39-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:40-Explanation of artificial sequence: Synthetic
DNA
5 SEQ ID NO:41-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:42-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:43-Explanation of artificial sequence: Synthetic
10 DNA
SEQ ID NO:44-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:45-Explanation of artificial sequence: Synthetic
DNA
15 SEQ ID NO:46-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:47-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:48-Explanation of artificial sequence: Synthetic
20 DNA
SEQ ID NO:49-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:50-Explanation of artificial sequence: Synthetic
DNA
25 SEQ ID NO:51-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:52-Explanation of artificial sequence: Synthetic
DNA
SEQ ID NO:53-Explanation of artificial sequence: Amino acid
30 sequence obtained by synthetic DNA
SEQ ID NO:54-Explanation of artificial sequence: Amino acid
sequence obtained by synthetic DNA
SEQ ID NO:57-Explanation of artificial sequence: Amino acid
sequence obtained by synthetic DNA

CLAIMS

1. A derivative of an antibody, comprising a monoclonal antibody or the antibody fragment thereof which specifically reacts with ganglioside GD3 which is conjugated with a radioisotope, a protein or a low molecular weight agent.

2. The derivative of an antibody according to claim 1, wherein the monoclonal antibody which specifically reacts with ganglioside GD3 is an antibody selected from an antibody produced by a hybridoma, a humanized antibody and a human antibody.

3. The derivative of an antibody according to claim 1 or 2, wherein the monoclonal antibody comprises CDR1, CDR2 and CDR3 of H chain V region having the amino acid sequences represented by SEQ ID NOS:3, 4 and 5, respectively.

4. The derivative of an antibody according to claim 1 or 2, wherein the monoclonal antibody comprises CDR1, CDR2 and CDR3 of L chain V region having the amino acid sequences represented by SEQ ID NOS:6, 7 and 8, respectively.

5. The derivative of an antibody according to claim 1 or 2, wherein the monoclonal antibody comprises:

CDR1, CDR2 and CDR3 of a heavy chain (H chain) variable region (V region) having the amino acid sequences represented by SEQ ID NOS:3, 4 and 5, respectively; and

CDR1, CDR2 and CDR3 of a light chain (L chain) V region having the amino acid sequences represented by SEQ ID NOS:6, 7 and 8, respectively.

6. The derivative of an antibody according to claim 2, wherein the antibody produced by a hybridoma is KM641 (FERM BP-3116).

7. The derivative of an antibody according to claim 2, wherein the humanized antibody is a human chimeric antibody or a human CDR-grafted antibody.

8. The derivative of a human chimeric antibody according to claim 7, wherein the human chimeric antibody comprises an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3 produced by a hybridoma.

9. The derivative of a human chimeric antibody according to claim 7, wherein the human chimeric antibody comprises:

an H chain V region and an L chain V region of a monoclonal antibody produced against ganglioside GD3 by a hybridoma; and

an H chain constant region (C region) and an L chain C region of a human antibody.

10. The derivative of a human chimeric antibody according to claim 8 or 9, wherein the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55.

11. The derivative of a human chimeric antibody according to claim 8 or 9, wherein the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

12. The derivative of a human chimeric antibody according to claim 8 or 9, wherein

the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55; and

the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

13. The derivative of a human chimeric antibody KM871 according to claim 8 or 9, wherein

the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55; and

the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

14. The derivative of a human CDR-grafted antibody according to claim 7, wherein the human CDR-grafted antibody comprises CDR of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3.

15. The derivative of a human CDR-grafted antibody according to claim 7, wherein the human CDR-grafted antibody comprises:

CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3; and

framework regions (FRs) of an H chain V region and an L chain V region of a human antibody.

16. The derivative of an antibody according to claim 7, wherein the human CDR-grafted antibody comprises:

CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3;

FRs of an H chain V region and an L chain V region of a human antibody; and

an H chain C region and an L chain C region of a human antibody.

17. The derivative of a human CDR-grafted antibody according to any one of claims 14 to 16, wherein the antibody comprises CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

18. The derivative of a human CDR-grafted antibody according to any one of claims 14 to 16, wherein the antibody comprises CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

19. The derivative of a human CDR-grafted antibody according to any one of claims 14 to 16, wherein the antibody comprises:

CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively; and

CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

20. The derivative of a human CDR-grafted antibody according to any one of claims 14 to 16, wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9.

21. The derivative of a human CDR-grafted antibody according to any one of claims 14 to 16, wherein the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

22. The derivative of a human CDR-grafted antibody according to any one of claims 14 to 16, wherein the H chain V region and the L chain V region of the antibody

comprises the amino acid sequences represented by SEQ ID NO:9 and SEQ ID NO:54, respectively.

23. The derivative of a human CDR-grafted antibody KM8871 according to any one of claims 14 to 16, wherein

the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and

the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

24. The derivative of the antibody fragment according to claim 1, wherein the antibody fragment is an antibody fragment selected from Fab, Fab', F(ab')₂, a single chain antibody (scFv), a disulfide stabilized V region fragment (dsFv) and a peptide comprising CDR.

25. The derivative of the antibody fragment according to claim 1 or 24, wherein the antibody fragment comprises amino acid sequences of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3 produced by a hybridoma.

26. The derivative of the antibody fragment according to claim 1 or 24, wherein the antibody fragment comprises an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:55.

27. The derivative of the antibody fragment according to claim 1 or 24, wherein the antibody fragment comprises an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:56.

28. The derivative of the antibody fragment according to claim 1 or 24, wherein the antibody fragment comprises:

an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:55; and

an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:56.

29. The derivative of the antibody fragment according to claim 1 or 24, wherein the antibody fragment comprises amino acid sequences of an H chain V region and an L chain V region of a human CDR-grafted antibody against ganglioside GD3.

30. The derivative of the antibody fragment according to claim 1 or 24, wherein the antibody fragment comprises an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:9.

31. The derivative of the antibody fragment according to claim 1 or 24, wherein the antibody fragment comprises an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:54.

32. The derivative of the antibody fragment according to claim 1 or 24, wherein the antibody fragment comprises:

an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:9; and

an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:54.

33. The derivative of an antibody fragment according to claim 1 or 24, wherein the antibody fragment comprises CDR1, CDR2 and CDR3 of an H chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

34. The derivative of the antibody fragment according to claim 1 or 24, wherein the antibody fragment comprises CDR1, CDR2 and CDR3 of an L chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

35. The derivative of the antibody fragment according to claim 1 or 24, wherein the antibody fragment comprises:

CDR1, CDR2 and CDR3 of an H chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5; and

CDR1, CDR2 and CDR3 of an L chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8.

36. The derivative of a monoclonal antibody or the antibody fragment thereof according to any one of claim 1 to 35, wherein the protein is a cytokine.

37. The derivative of a monoclonal antibody or the antibody fragment thereof according to claim 36, wherein the cytokine is human interleukin-2 (hIL-2).

38. The derivative of an antibody according to claim 37, wherein the derivative of an antibody comprises a human chimeric antibody KM871 and hIL-2.

39. The derivative of an antibody according to claim 38, wherein the antibody conjugated with hIL-2 comprises:

an H chain V region having the amino acid sequence represented by SEQ ID NO:57; and

an L chain V region having the amino acid sequence represented by SEQ ID NO:56.

40. The derivative of an antibody according to claim 37, wherein the derivative of an antibody comprises a human CDR-grafted antibody KM8871 and hIL-2.

41. The derivative of an antibody according to claim 1, wherein the antibody conjugated with hIL-2 comprises:

an H chain V region having the amino acid sequence represented by SEQ ID NO:53; and

an L chain V region having the amino acid sequence represented by SEQ ID NO:54.

42. A DNA which encodes the derivative of a monoclonal antibody or the derivative of the antibody fragment thereof which specifically reacts with ganglioside GD3 according to any one of claims 1 to 41.

43. A recombinant vector comprising the DNA according to claim 42.

44. A transformant which is obtained by introducing the recombinant vector according to claim 43 into a host cell.

45. A transformant KM871hIL2 (FERM BP-6918) which produces the antibody according to claim 38.

46. A transformant KM8871hIL2 (FERM BP-6791) which produces the antibody according to claim 40.

47. A process for producing an antibody, which comprises:

culturing the transformant according to any one of claims 44 to 46 in a culture medium to produce and

accumulate the derivative of a monoclonal antibody or the derivative of the antibody fragment thereof according to any one of claims 1 to 41 in the culture; and

recovering the derivative of the antibody or the derivative of the antibody fragment thereof from the culture.

48. A human CDR-grafted antibody or the antibody fragment thereof which specifically reacts with ganglioside GD3.

49. The human CDR-grafted antibody or the antibody fragment thereof according to claim 48, wherein the human CDR-grafted antibody comprises CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3.

50. The human CDR-grafted antibody or the antibody fragment thereof according to claim 48, wherein the human CDR-grafted antibody comprises:

CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3; and

FRs of an H chain V region and an L chain V region of a human antibody.

51. The human CDR-grafted antibody or the antibody fragment thereof according to claim 48, wherein the human CDR-grafted antibody comprises:

CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3;

FRs of an H chain V region and an L chain V region of a human antibody; and

an H chain C region and an L chain C region of a human antibody.

52. The human CDR-grafted antibody or the antibody fragment thereof according to any one of claims 49 to 51, wherein the antibody comprises CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

53. The human CDR-grafted antibody or the antibody fragment thereof according to any one of claims 49 to 51, wherein the antibody comprises CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

54. The human CDR-grafted antibody or the antibody fragment thereof according to any one of claims 49 to 51, wherein the antibody comprises:

CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively; and

CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8.

55. The human CDR-grafted antibody or the antibody fragment thereof according to any one of claims 49 to 51, wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9.

56. The human CDR-grafted antibody or the antibody fragment thereof according to any one of claims 49 to 51, wherein the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

57. The human CDR-grafted antibody or the antibody fragment thereof according to any one of claims 49 to 51, wherein

the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and

the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

58. The human CDR-grafted antibody KM8871 or the antibody fragment thereof according to any one of claims 49 to 51, wherein

the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and

the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

59. A DNA which encodes the human CDR-grafted antibody or the antibody fragment thereof which specifically reacts with ganglioside GD3 according to any one of claims 48 to 58.

60. A recombinant vector comprising the DNA according to claim 59.

61. A transformant which is obtained by introducing the recombinant vector according to claim 60 into a host cell.

62. A transformant KM8871 (FERM BP-6790) which produces the human CDR-grafted antibody according to claim 58.

63. A process for producing an antibody, which comprises:

culturing the transformant according to claim 61 or 62 in a culture medium to produce and accumulate the human CDR-grafted antibody or the antibody fragment thereof according to any one of claims 48 to 58 in the culture; and

recovering the antibody or the antibody fragment thereof from the culture.

64. A medicament comprising at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to claims 1 to 41 and the human CDR-grafted antibody and the antibody fragment thereof according to any one of claims 48 to 58.

65. A therapeutic agent for cancers, comprising, as an active ingredient, at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to any one of claims 1 to 41 and the human CDR-grafted antibody and the antibody fragment thereof according to any one of claims 48 to 58.

66. A diagnostic agent for cancers, comprising, as an active ingredient, at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to any one of claims 1 to 41 and the human CDR-grafted antibody and the antibody fragment thereof according to any one of claims 48 to 58.


ABSTRACT

The present invention relates to a human CDR-grafted antibody against ganglioside GD3 (hereinafter referred to "GD3"), derivatives of an anti-GD3 antibody and cytokine, and use for treatment and diagnosis of the
5 antibody and the derivatives.

1 / 41

Fig.

1




SEQ ID NO:

配列番号11~16

2 / 41

Fig.


2



3 / 41

Fig. 3


3



4 / 41

Fig.


4



5 / 41

Fig.


5



6 / 41


Fig.

6




7 / 41

Fig.
7



8 / 41


Fig.
8



9 / 41

Fig.


9



10 / 41

Fig.


10



11 / 41

Fig.


11



12 / 41

Fig


1 2




13 / 41

Fig.

13




14



14 / 41

Fig.


15



15 / 41


Fig. 16

16



16 / 41



Fig. 17



17 / 41



Fig.

18




18 / 41

fig
19



19 / 41


fig.
20



20 / 41

Fig.


2 1



21 / 41

Fig.


22



22 / 41

Fig.

2 3



23 / 41

Fig.

24





Fig.


25



24 / 41

Fig.

26



25 / 41

Fig.

27

cell
G361細胞






Fig.
28




27 / 41

fig.

29

Cytotoxic activity
細胞障害活性(%)



28 / 41

Fig.
30






Fig.

3 1




3 2



30 / 41

Fig.

33





結合活性(OD415)

Binding activity



31 / 41

Fig.
3 4

Cell
G361細胞




Cell
SK-MEL-28細胞



32 / 41

Fig.

35



33 / 41

Fig

☒ 3 6






Fig.
37



35 / 41


Fig.

 38




B16 · F10 (親株)

Parent strain



B16 - 29-10




B16 - 8-3

36 / 41

Fig.

39

Number of metastatic foci
転移巣の数(個)



バーは各群の平均値

Bar: Average value of each group

No administration


KM871
-hIL-2

KM871

37 / 41

Fig.


40



No administration

Fig.

41



No administration

○ KM871-hIL-2


● KM871+hIL-2 併用投与

Combined administration

38 / 41


Fig. 42

Tumor volume of each group / tumor volume of non-administered group



Days after administration started Fig. 43

v/v0 of each group / v/v0 of non-administered group




Days after administration started

39 / 41

Fig.
44

v/v0 of each group
/ v/v0 of non-administered group
各群のv/v0 / 非投与群のv/v0




投与開始後の日数
Days after administration started

-コントロール Control
- ▲ hIL-2
- KM871
- KM871+KM871-hIL-2 15µg 併用 Combined use
- ◇ KM871+KM871-hIL-2 80µg 併用 Combined use

Fig

4 5



KM871 concentration used for assay

4 6

Fig.




Fig.

47





Fig.

48



縦軸: 転移果の数 バー: 各群の平均値
 グラフ中の数値: KM871-hIL-2投与による転移阻害率

Ordinate: number of metastatic foci
 Bar: average of each group
 Value in the graph: metastasis inhibition ratio by KM871-hIL-2 administration

Case No. _____

Nixon & Vanderhye P.C. (12/97)

RULE 63 (37 C.F.R. 1.63)
DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION
IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

As a below named inventor, I hereby declare that my residence, post office address and citizenship are as stated below next to my name, and I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: **COMPLEMENTARITY DETERMINING REGION-GRAFTED ANTIBODY AGAINST GANGLIOSIDE GD3 AND DERIVATIVE OF ANTIBODY AGAINST GANGLIOSIDE GD3**

the specification of which (check applicable box(es)):

☐ is attached hereto
☐ was filed on _____ as U.S. Application Serial No. _____
☒ was filed as PCT International application No. PCT/JP00/06774 on September 29, 2000
and (if applicable to U.S. or PCT application) was amended on _____

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with 37 C.F.R. 1.58. I hereby claim foreign priority benefits under 35 U.S.C. 119/365 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed or, if no priority is claimed, before the filing date of this application:

Priority Foreign Application(s):

Application Number	Country	Day/Month/Year Filed
P. Hei. 11-278291	Japan	30/September/1999
P. 2000-105088	Japan	6/April/2000

I hereby claim the benefit under 35 U.S.C. §119(e) of any United States provisional application(s) listed below:

Application Number	Date/Month/Year Filed
--------------------	-----------------------

I hereby claim the benefit under 35 U.S.C. 120/365 of all prior United States and PCT International applications listed above or below and, insofar as the subject matter of each of the claims of this application is not disclosed in such prior applications in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose material information as defined in 37 C.F.R. 1.56 which occurred between the filing date of the prior applications and the national or PCT international filing date of this application:

Prior U.S./PCT Application(s):

Application Serial No.	Day/Month/Year Filed	Status: patented pending, abandoned
------------------------	----------------------	--

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon. And I hereby appoint **NIXON & VANDERHYE P.C., 1100 North Glebe Rd., 8th Floor, Arlington, VA 22201-4714, telephone number (703) 816-4000** (to whom all communications are to be directed), and the following attorneys thereof (of the same address) individually and collectively my attorneys to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith and with the resulting patent: Arthur R. Crawford, 25327; Larry S. Nixon, 25640; Robert A. Vanderhye, 27076; James T. Hosmer, 30184; Robert W. Farris, 31352; Richard G. Besha, 22770; Mark E. Nusbaum, 32348; Michael J. Keenan, 32106; Bryan H. Davidson, 30251; Stanley C. Spooner, 27393; Leonard C. Mitchard, 29009; Duane M. Byers, 33363; Jeffry H. Nelson, 30481; John R. Lastova, 33149; H. Warren Burnam, Jr. 29366; Thomas E. Byrne, 32205; Mary J. Wilson, 32955; J. Scott Davidson, 33489; Alan M. Kagen, 36178; William J. Griffin, 31260; Robert A. Molan, 29834; B. J. Sadoff, 36663; James D. Berquist, 34776; Updeep S. Gill, 37334.

1. Inventor's Signature: Nobuo Hanai Date: March 25, 2002

Inventor: Nobuo HANAI Japan
(first) (last) (citizenship)
Residence: (city) Machida-shi, Tokyo JP (state/country) Japan
Post Office Address: c/o Tokyo Research Laboratories, KYOWA HAKKO KOGYO CO., LTD., 6-6, Asahi-machi 3-chome,
(Zip Code) Machida-shi, Tokyo 194-8533 Japan

2. Inventor's Signature: Kenya Shitara Date: March 25, 2002

Inventor: Kenya SHITARA Japan
(first) (last) (citizenship)
Residence: (city) Machida-shi, Tokyo JP (state/country) Japan
Post Office Address: c/o Tokyo Research Laboratories, KYOWA HAKKO KOGYO CO., LTD., 6-6, Asahi-machi 3-chome,
(Zip Code) Machida-shi, Tokyo 194-8533 Japan

3. Inventor's Signature: Kazuyasu Nakamura Date: March 25, 2002

Inventor: Kazuyasu NAKAMURA Japan
(first) (last) (citizenship)
Residence: (city) Machida-shi, Tokyo JP (state/country) Japan
Post Office Address: c/o Tokyo Research Laboratories, KYOWA HAKKO KOGYO CO., LTD., 6-6, Asahi-machi 3-chome,
(Zip Code) Machida-shi, Tokyo 194-8533 Japan

FOR ADDITIONAL INVENTORS, check box ☒ and attach sheet with same information and signature and date for each.

Case No. _____

Nixon & Vandemye P.C. (12/97)

RULE 63 (37 C.F.R. 1.63)
DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION
IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

4.	Inventor's Signature: _____ Inventor: _____	<u>Rinpei Niwa</u> <div style="display: flex; justify-content: space-between; font-size: small;"> (first) MI (last) </div>	Date: <u>March 25, 2002</u> <div style="display: flex; justify-content: space-between; font-size: small;"> NIWA Japan </div> <div style="display: flex; justify-content: space-between; font-size: small;"> (first) (last) (citizenship) </div>
	Residence: (city) _____ Post Office Address: _____ (Zip Code) _____	<u>Machida-shi, Tokyo</u> <u>Japan</u> <u>c/o Tokyo Research Laboratories, KYOWA HAKKO KOGYO CO., LTD., 6-6, Asahi-machi 3-chome.</u> <u>Machida-shi, Tokyo 194-8533 Japan</u>	
5.	Inventor's Signature: _____ Inventor: _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (first) MI (last) </div>	Date: _____ <div style="display: flex; justify-content: space-between; font-size: small;"> </div> <div style="display: flex; justify-content: space-between; font-size: small;"> (citizenship) </div>
	Residence: (city) _____ Post Office Address: _____ (Zip Code) _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (state/country) </div>	
6.	Inventor's Signature: _____ Inventor: _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (first) MI (last) </div>	Date: _____ <div style="display: flex; justify-content: space-between; font-size: small;"> </div> <div style="display: flex; justify-content: space-between; font-size: small;"> (citizenship) </div>
	Residence: (city) _____ Post Office Address: _____ (Zip Code) _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (state/country) </div>	
7.	Inventor's Signature: _____ Inventor: _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (first) MI (last) </div>	Date: _____ <div style="display: flex; justify-content: space-between; font-size: small;"> </div> <div style="display: flex; justify-content: space-between; font-size: small;"> (citizenship) </div>
	Residence: (city) _____ Post Office Address: _____ (Zip Code) _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (state/country) </div>	
8.	Inventor's Signature: _____ Inventor: _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (first) MI (last) </div>	Date: _____ <div style="display: flex; justify-content: space-between; font-size: small;"> </div> <div style="display: flex; justify-content: space-between; font-size: small;"> (citizenship) </div>
	Residence: (city) _____ Post Office Address: _____ (Zip Code) _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (state/country) </div>	
9.	Inventor's Signature: _____ Inventor: _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (first) MI (last) </div>	Date: _____ <div style="display: flex; justify-content: space-between; font-size: small;"> </div> <div style="display: flex; justify-content: space-between; font-size: small;"> (citizenship) </div>
	Residence: (city) _____ Post Office Address: _____ (Zip Code) _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (state/country) </div>	
10.	Inventor's Signature: _____ Inventor: _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (first) MI (last) </div>	Date: _____ <div style="display: flex; justify-content: space-between; font-size: small;"> </div> <div style="display: flex; justify-content: space-between; font-size: small;"> (citizenship) </div>
	Residence: (city) _____ Post Office Address: _____ (Zip Code) _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (state/country) </div>	
11.	Inventor's Signature: _____ Inventor: _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (first) MI (last) </div>	Date: _____ <div style="display: flex; justify-content: space-between; font-size: small;"> </div> <div style="display: flex; justify-content: space-between; font-size: small;"> (citizenship) </div>
	Residence: (city) _____ Post Office Address: _____ (Zip Code) _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (state/country) </div>	
12.	Inventor's Signature: _____ Inventor: _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (first) MI (last) </div>	Date: _____ <div style="display: flex; justify-content: space-between; font-size: small;"> </div> <div style="display: flex; justify-content: space-between; font-size: small;"> (citizenship) </div>
	Residence: (city) _____ Post Office Address: _____ (Zip Code) _____	<div style="display: flex; justify-content: space-between; font-size: small;"> (state/country) </div>	

107089500

JC13 Rec'd PCT/PTO 29 MAR 2002

SEQUENCE LISTING

<110> KYOWA HAKKO KOGYO CO., LTD

<120> Humanized anti-GD3 antibody and it's cytokine conjugate

<130>11239WO1

<140>

<141>

<150>H11-278291

<151>1999-09-30

<160> 57

<170> PatentIn Ver. 2.0

<210> 1

<211> 138

<212> PRT

<213> Mus musculus

<400> 1

Met Glu Phe Gly Leu Ser Trp Leu Phe Leu Val Leu Val Phe Lys Gly
-19 -15 -10 -5

Val Gln Cys Glu Val Thr Leu Val Glu Ser Gly Gly Asp Phe Val Lys
-1 1 5 10

Pro Gly Gly Ser Leu Lys Val Ser Cys Ala Ala Ser Gly Phe Ala Phe
15 20 25

Ser His Tyr Ala Met Ser Trp Val Arg Gln Thr Pro Ala Lys Arg Leu
30 35 40 45

Glu Trp Val Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser
50 55 60

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
65 70 75

Thr Leu Tyr Leu Gln Met Arg Ser Leu Arg Ser Glu Asp Ser Ala Met
80 85 90

Tyr Phe Cys Thr Arg Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser Trp
95 100 105

Gly Gln Gly Thr Thr Leu Thr Val Ser Ser
110 115

<210> 2

<211> 128
<212> PRT
<213> Mus musculus

<400> 2
Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Leu Cys Phe Gln
-20 -15 -10 -5
Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Thr Ala Ser Ser Leu Pro
-1 1 5 10
Ala Ser Leu Gly Asp Arg Val Thr Ile Ser Cys Ser Ala Ser Gln Asp
15 20 25
Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val
30 35 40
Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
45 50 55 60
Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser
65 70 75
Asn Leu Glu Pro Glu Asp Ile Ala Thr Tyr Phe Cys His Gln Tyr Ser
80 85 90
Lys Leu Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
95 100 105

<210> 3
<211> 5
<212> PRT
<213> Mus musculus

<400> 3
His Tyr Ala Met Ser
1 5

<210> 4
<211> 17
<212> PRT
<213> Mus musculus

<400> 4
Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val Lys Gly
1 5 10 15

<210> 5
<211> 10
<212> PRT
<213> Mus musculus

<400> 5

Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser
1 5 10

<210> 6
<211> 11
<212> PRT
<213> Mus musculus

<400> 6
Ser Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn
1 5 10

<210> 7
<211> 7
<212> PRT
<213> Mus musculus

<400> 7
Tyr Ser Ser Asn Leu His Ser
1 5

<210> 8
<211> 9
<212> PRT
<213> Mus musculus

<400> 8
His Gln Tyr Ser Lys Leu Pro Trp Thr
1 5

<210> 9
<211> 119
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic protein

<400> 9
Glu Val Gln Leu Val Glu Ser Gly Gly Asp Phe Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Val Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser His Tyr
20 25 30

Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr

<210> 12
<211> 97
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 12
aaagcgaatc cagaggctgc acaggagact ctcagagacc cccccggctg tacaaagtct 60
ccccagact ccaccagctg cacctcacac tgaacac 97

<210> 13
<211> 98
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 13
gcagcctctg gattcgcttt cagtcattat gccatgtctt ggggccgcca ggctccaggg 60
aaggggctgg agtgggtggc ttatattagt agtgggtgg 98

<210> 14
<211> 97
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 14
gatacagcgt gttcttggag ttatctctgg agatgggtgaa tctgcccttt acactgtctg 60
aatagtaggt gccactacca ccactactaa tataagc 97

<210> 15
<211> 98
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 15
ctccaagaac acgctgtatc tgcaaattgcg cagcctgaga gctgaggact cggctgtgta 60
tttctgtaca agagttaaac tgggaaccta ctactttg 98

<210> 16
<211> 92
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 16

gttttcccag tcacgacggg cccttggtgg aggctgagga gacggtgagc agggttccct 60
ggccccagga gtcaaagtag taggttccca gt 92

<210> 17

<211> 94

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 17

caggaaacag ctatgacgaa ttccaccatg atgtcctctg ctcagttcct tggctctctg 60
ttgctctggt ttcaaggtag cagatgtgac atcc 94

<210> 18

<211> 87

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 18

gcactacaag tgatggtgac tctgtctcct acagatgcag acagggagga tggagactgg 60
gtcatctgga tgtcacatct ggtacct 87

<210> 19

<211> 89

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 19

gtcaccatca cttgtagtgc aagtcaggac attagtaatt atttaaactg gtatcagcag 60
aaaccaggga aagcccctaa gctcctgat 89

<210> 20

<211> 89

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 20
taatctgtcc cagatccacc gccgctgaac cttgatggga ccccgagtg taaatttgat 60
gagtaaaaga tcaggagctt aggggcttt 89

<210> 21
<211> 92
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 21
ggtggatctg ggacagatta tactctcacc atcagcagcc tgcagcctga agattttgca 60
acttattact gtcacagta tagtaagctt cc 92

<210> 22
<211> 80
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 22
gttttccag tcacgaccgt acgtttaatc tctaccttgg tcccctggcc gaacgtccac 60
ggaagcttac tatactgatg 80

<210> 23
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 23
agcttccatg gacgttcggt ggaggcacca agctggaaat caaac 45

<210> 24
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 24
gtacgtttga tttccagctt ggtgcctcca ccgaacgtcc atgga 45

<210> 25
 <211> 29
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<400> 25
 gatgcaggca gggaggatgc agtctgggt 29

<210> 26
 <211> 29
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<400> 26
 acccagactg catcctccct gcctgcatc 29

<210> 27
 <211> 384
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<220>
 <221> CDS
 <222> (1)..(384)

<400> 27
 atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
 Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
 -20 -15 -10 -5

ggg acc aga tgt gac atc cag atg acc cag act gca tcc tcc ctg cct 96
 Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Thr Ala Ser Ser Leu Pro
 -1 1 5 10

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
 Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
 15 20 25

att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc cct 192
 Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro
 30 35 40

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240
 Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
 45 50 55 60

agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288
 Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser
 65 70 75

agc ctg cag cct gaa gat ttt gca act tat tac tgt cat cag tat agt 336
 Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys His Gln Tyr Ser
 80 85 90

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384
 Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
 95 100 105

<210> 28
 <211> 26
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<400> 28
 actgatgaca gaaataagtt gcaaaa 26

<210> 29
 <211> 26
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<400> 29
 ttttgcaact tatttctgtc atcagt 26

<210> 30
 <211> 384
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<220>
 <221> CDS
 <222> (1)..(384)

<400> 30
 atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
 Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Leu Cys Phe Gln
 -20 -15 -10 -5

ggg acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<220>

<221> CDS

<222> (1)..(384)

<400> 33

atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa	48
Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Leu Cys Phe Gln	
-20 -15 -10 -5	

ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct	96
Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser	
-1 1 5 10	

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac	144
Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp	
15 20 25	

att agt aat tat tta aac tgg tat cag cag aaa cca gac aaa gcc gtt	192
Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Asp Lys Ala Val	
30 35 40	

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca	240
Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser	
45 50 55 60	

agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc	288
Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser	
65 70 75	

agc ctg cag cct gaa gat ttt gca act tat tac tgt cat cag tat agt	336
Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys His Gln Tyr Ser	
80 85 90	

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt	384
Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg	
95 100 105	

<210> 34

<211> 48

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 34

gttgcgatat cttcaggctg cagattgctg atggtgagac tataatct	48
--	----

<210> 35

<211> 48
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 35
agattatagt ctcaccatca gcaatctgca gcctgaagat atcgcaac 48

<210> 36
<211> 384
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<220>
<221> CDS
<222> (1)..(384)

<400> 36
atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
-20 -15 -10 -5

ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96
Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
-1 1 5 10

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
15 20 25

att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc cct 192
Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro
30 35 40

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240
Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
45 50 55 60

agg ttc agc ggc ggt gga tct ggg aca gat tat agt ctc acc atc agc 288
Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser
65 70 75

aat ctg cag cct gaa gat atc gca act tat tac tgt cat cag tat agt 336
Asn Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys His Gln Tyr Ser
80 85 90

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384
Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
95 100 105

<210> 37
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<400> 37
 ttcaggctgc agattgctga tgggtg 25

<210> 38
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<400> 38
 caccatcagc aatctgcagc ctgaa 25

<210> 39
 <211> 384
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<220>
 <221> CDS
 <222> (1)..(384)

<400> 39
 atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
 Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
 -20 -15 -10 -5

ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96
 Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
 -1 1 5 10

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
 Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
 15 20 25

att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc cct 192
 Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro
 30 35 40

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240
 Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
 45 50 55 60

agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288
 Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser
 65 70 75

aat ctg cag cct gaa gat ttt gca act tat tac tgt cat cag tat agt 336
 Asn Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys His Gln Tyr Ser
 80 85 90

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384
 Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
 95 100 105

<210> 40
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<400> 40
 gacagaaata agttgcgata tcttcaggct 30

<210> 41
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<400> 41
 agcctgaaga tatcgcaact tatttctgtc 30

<210> 42
 <211> 384
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<220>
 <221> CDS
 <222> (1)..(384)

<400> 42
 atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
 Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Leu Cys Phe Gln
 -20 -15 -10 -5

ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 45
taatctgtcc cagatccacc gccgctgaac cttgatggga cccccgagtg taaatttgat 60
gagtaaaaga tcaggagctt aacggcttt 89

<210> 46
<211> 92
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 46
ggtggatctg ggacagatta tactctcacc atcagcagcc tgcagcctga agattttgca 60
acttatttct gtcatcagta tagtaagctt cc 92

<210> 47
<211> 384
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<220>
<221> CDS
<222> (1)..(384)

<400> 47
atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
-20 -15 -10 -5
ggt acc aga tgt gac atc cag atg acc cag tct gca tcc tcc ctg cct 96
Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Ala Ser Ser Leu Pro
-1 1 5 10
gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
15 20 25
att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc gtt 192
Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Val
30 35 40
aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240
Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
45 50 55 60

agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288
 Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser
 65 70 75

agc ctg cag cct gaa gat ttt gca act tat ttc tgt cat cag tat agt 336
 Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Phe Cys His Gln Tyr Ser
 80 85 90

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384
 Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
 95 100 105

<210> 48

<211> 384

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<220>

<221> CDS

<222> (1)..(384)

<400> 48

atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
 Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Leu Cys Phe Gln
 -20 -15 -10 -5

ggc acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96
 Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
 -1 1 5 10

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
 Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
 15 20 25

att agt aat tat tta aac tgg tat cag cag aaa cca gac aaa gcc gtt 192
 Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Asp Lys Ala Val
 30 35 40

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240
 Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
 45 50 55 60

agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288
 Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser
 65 70 75

agc ctg cag cct gaa gat atc gca act tat ttc tgt cat cag tat agt 336
 Ser Leu Gln Pro Glu Asp Ile Ala Thr Tyr Phe Cys His Gln Tyr Ser
 80 85 90

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384

Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
95 100 105

<210> 49
<211> 76
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 49
catgcatgag gctctgcaca accactacac gcagaagagc ctctccctgt ctcccggggg 60
agaattcatt gatcag 76

<210> 50
<211> 85
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 50
gatcctgata aatgaattct cccccgggag acagggagag gctcttctgc gtgtagtggt 60
tgtgcagagc ctcatgcatg gggcc 85

<210> 51
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 51
gtctcccgga aaagcaccta ctagtagttc taaaaag 37

<210> 52
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 52
ccctgatcaa tgaattcaag tcagtgttga gatgatgc 38

<210> 53

<223> Description of Artificial Sequence:synthetic protein

Glu Val Gln Leu Val Glu Ser Gly Gly Asp Phe Val Gln Pro Gly Gly
1 5 10 15

Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Arg Ser Leu Arg Ala Glu Asp Ser Ala Val Tyr Phe Cys
85 90 95

Thr Arg Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser Trp Gly Gln Gly
100 105 110

Thr Leu Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
115 120 125

Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
130 135 140

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
145 150 155 160

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
165 170 175

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
180 185 190

Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro
195 200 205

Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys
210 215 220

Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro
225 230 235 240

Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser
245 250 255

Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp

260										265					270				
Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn				
		275					280					285							
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val				
		290					295					300							
Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu				
305					310					315			320						
Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys				
				325					330					335					
Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr				
				340					345					350					
Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr				
				355					360					365					
Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu				
						375					380								
Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu				
385					390					395					400				
Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys				
				405					410					415					
Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu				
				420					425					430					
Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly				
				435					440					445					
Lys	Ala	Pro	Thr	Ser	Ser	Ser	Thr	Lys	Lys	Thr	Gln	Leu	Gln	Leu	Glu				
						455					460								
His	Leu	Leu	Leu	Asp	Leu	Gln	Met	Ile	Leu	Asn	Gly	Ile	Asn	Asn	Tyr				
465					470					475					480				
Lys	Asn	Pro	Lys	Leu	Thr	Arg	Met	Leu	Thr	Phe	Lys	Phe	Tyr	Met	Pro				
				485					490					495					
Lys	Lys	Ala	Thr	Glu	Leu	Lys	His	Leu	Gln	Cys	Leu	Glu	Glu	Glu	Leu				
				500					505					510					
Lys	Pro	Leu	Glu	Glu	Val	Leu	Asn	Leu	Ala	Gln	Ser	Lys	Asn	Phe	His				
				515					520					525					
Leu	Arg	Pro	Arg	Asp	Leu	Ile	Ser	Asn	Ile	Asn	Val	Ile	Val	Leu	Glu				
						535					540								
Leu	Lys	Gly	Ser	Glu	Thr	Thr	Phe	Met	Cys	Glu	Tyr	Ala	Asp	Glu	Thr				
545					550					555					560				
Ala	Thr	Ile	Val	Glu	Phe	Leu	Asn	Arg	Trp	Ile	Thr	Phe	Cys	Gln	Ser				

Ile Ile Ser Thr Leu Thr
580

<210> 54
<211> 108
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic protein

<400> 54
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15
Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr
20 25 30
Leu Asn Trp Tyr Gln Gln Lys Pro Asp Lys Ala Val Lys Leu Leu Ile
35 40 45
Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60
Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80
Glu Asp Ile Ala Thr Tyr Phe Cys His Gln Tyr Ser Lys Leu Pro Trp
85 90 95
Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
100 105

<210> 55
<211> 119
<212> PRT
<213> Mus musculus

<400> 55
Glu Val Thr Leu Val Glu Ser Gly Gly Asp Phe Val Lys Pro Gly Gly
1 5 10 15
Ser Leu Lys Val Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser His Tyr
20 25 30
Ala Met Ser Trp Val Arg Gln Thr Pro Ala Lys Arg Leu Glu Trp Val
35 40 45
Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Arg Ser Leu Arg Ser Glu Asp Ser Ala Met Tyr Phe Cys
85 90 95

Thr Arg Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser Trp Gly Gln Gly
100 105 110

Thr Thr Leu Thr Val Ser Ser
115

<210> 56

<211> 108

<212> PRT

<213> Mus musculus

<400> 56

Asp Ile Gln Met Thr Gln Thr Ala Ser Ser Leu Pro Ala Ser Leu Gly
1 5 10 15

Asp Arg Val Thr Ile Ser Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr
20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val Lys Leu Leu Ile
35 40 45

Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Gly Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser Asn Leu Glu Pro
65 70 75 80

Glu Asp Ile Ala Thr Tyr Phe Cys His Gln Tyr Ser Lys Leu Pro Trp
85 90 95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
100 105

<210> 57

<211> 582

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic protein

<400> 57

Glu Val Thr Leu Val Glu Ser Gly Gly Asp Phe Val Lys Pro Gly Gly
1 5 10 15

Ser Leu Lys Val Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser His Tyr
20 25 30

Ala Met Ser Trp Val Arg Gln Thr Pro Ala Lys Arg Leu Glu Trp Val
35 40 45

Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val

50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Arg Ser Leu Arg Ser Glu Asp Ser Ala Met Tyr Phe Cys
 85 90 95
 Thr Arg Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser Trp Gly Gln Gly
 100 105 110
 Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
 115 120 125
 Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
 130 135 140
 Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
 145 150 155 160
 Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
 165 170 175
 Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
 180 185 190
 Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro
 195 200 205
 Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys
 210 215 220
 Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro
 225 230 235 240
 Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser
 245 250 255
 Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp
 260 265 270
 Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn
 275 280 285
 Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val
 290 295 300
 Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu
 305 310 315 320
 Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys
 325 330 335
 Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr
 340 345 350
 Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr

355	360	365
Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu 370 375 380		
Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu 385 390 395 400		
Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys 405 410 415		
Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu 420 425 430		
Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly 435 440 445		
Lys Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu 450 455 460		
His Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr 465 470 475 480		
Lys Asn Pro Lys Leu Thr Arg Met Leu Thr Phe Lys Phe Tyr Met Pro 485 490 495		
Lys Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu 500 505 510		
Lys Pro Leu Glu Glu Val Leu Asn Leu Ala Gln Ser Lys Asn Phe His 515 520 525		
Leu Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu 530 535 540		
Leu Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr 545 550 555 560		
Ala Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser 565 570 575		
Ile Ile Ser Thr Leu Thr 580		

SEQUENCE LISTING

<110> KYOWA HAKKO KOGYO CO., LTD

<120> Humanized anti-GD3 antibody and it's cytokine conjugate

<130>11239W01

<140>

<141>

<150>H11-278291

<151>1999-09-30

<160> 57

<170> PatentIn Ver. 2.0

<210> 1

<211> 138

<212> PRT

<213> Mus musculus

<400> 1

Met Glu Phe Gly Leu Ser Trp Leu Phe Leu Val Leu Val Phe Lys Gly

-19

-15

-10

-5

Val Gln Cys Glu Val Thr Leu Val Glu Ser Gly Gly Asp Phe Val Lys

2 / 42

-1 1

5

10

Pro Gly Gly Ser Leu Lys Val Ser Cys Ala Ala Ser Gly Phe Ala Phe

15

20

25

Ser His Tyr Ala Met Ser Trp Val Arg Gln Thr Pro Ala Lys Arg Leu

30

35

40

45

Glu Trp Val Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser

50

55

60

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn

65

70

75

Thr Leu Tyr Leu Gln Met Arg Ser Leu Arg Ser Glu Asp Ser Ala Met

80

85

90

Tyr Phe Cys Thr Arg Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser Trp

95

100

105

Gly Gln Gly Thr Thr Leu Thr Val Ser Ser

110

115

<210> 2

<211> 128

<212> PRT

<213> Mus musculus

<400> 2

3 / 42

Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
-20 -15 -10 -5

Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Thr Ala Ser Ser Leu Pro
-1 1 5 10

Ala Ser Leu Gly Asp Arg Val Thr Ile Ser Cys Ser Ala Ser Gln Asp
15 20 25

Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val
30 35 40

Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
45 50 55 60

Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser
65 70 75

Asn Leu Glu Pro Glu Asp Ile Ala Thr Tyr Phe Cys His Gln Tyr Ser
80 85 90

Lys Leu Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
95 100 105

<210> 3

<211> 5

<212> PRT

<213> Mus musculus

<400> 3

His Tyr Ala Met Ser

1 5

<210> 4

<211> 17

<212> PRT

<213> Mus musculus

<400> 4

Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val Lys Gly

1 5 10 15

<210> 5

<211> 10

<212> PRT

<213> Mus musculus

<400> 5

Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser

1 5 10

<210> 6

<211> 11

<212> PRT

<213> Mus musculus

<400> 6

Ser Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn

1 5 10

<210> 7

<211> 7

<212> PRT

<213> Mus musculus

<400> 7

Tyr Ser Ser Asn Leu His Ser

1

5

<210> 8

<211> 9

<212> PRT

<213> Mus musculus

<400> 8

His Gln Tyr Ser Lys Leu Pro Trp Thr

1

5

<210> 9

<211> 119

<212> PRT

<213> Artificial Sequence

<400> 9

Glu Val Gln Leu Val Glu Ser Gly Gly Asp Phe Val Gln Pro Gly Gly

1

5

10

15

Ser Leu Arg Val Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser His Tyr

6 / 42

20

25

30

Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

35

40

45

Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val

50

55

60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr

65

70

75

80

Leu Gln Met Arg Ser Leu Arg Ala Glu Asp Ser Ala Val Tyr Phe Cys

85

90

95

Thr Arg Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser Trp Gly Gln Gly

100

105

110

Thr Leu Leu Thr Val Ser Ser

115

<210> 10

<211> 108

<212> PRT

<213> Artificial Sequence

<400> 10

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly

1

5

10

15

7 / 42

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr
20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys His Gln Tyr Ser Lys Leu Pro Trp
85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
100 105

<210> 11

<211> 95

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 11

caggaaacag ctatgaacgc gcgccacca tggagtttgg gctcagctgg ctttttcttg 60
tccttgtttt caaagggtgtt cagtgtgagg tgcag 95

<210> 12

<211> 97

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 12

aaagcgaatc cagaggctgc acaggagact ctcagagacc cccccggctg tacaaagtct 60
ccccagact ccaccagctg cacctcacac tgaacac 97

<210> 13

<211> 98

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 13

gcagcctctg gattogettt cagtcattat gccatgtctt gggtccgccg ggtccagg 60
aaggggctgg agtgggtggc ttatattagt agtgggtg 98

<210> 14

<211> 97

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 14

gatacagegt gttcttggag ttatctctgg agatggtgaa tetgcccctt acactgtctg 60
aatagtaggt gccactacca ccactactaa tataagc 97

<210> 15

<211> 98

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 15

ctccaagaac acgctgtatc tgcaaatgcg cagcctgaga gctgaggact cggtgtgtga 60
tttctgtaca agagttaaac tgggaaccta ctactttg 98

<210> 16

<211> 92

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 16

gttttccag tcacgacggg cccttggtgg aggtgagga gacggtgagc aggttccct 60
ggccccagga gtcaaagtag taggttccca gt 92

<210> 17

<211> 94

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 17

caggaaacag ctatgacgaa ttccaccatg atgtectctg ctcagttcct tggctctctg 60
ttgctctgtt ttcaaggtag cagatgtgac atcc 94

<210> 18

<211> 87

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 18

gcactacaag tgatggtagac tctgtctcct acagatgcag acagggagga tggagactgg 60
gtcatctgga tgccacatct ggtacct 87

<210> 19

<211> 89

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 19

gtcaccatca cttgtagtgc aagtcaggac attagtaatt atttaaactg gtagcagcag 60
aaaccaggga aagcccctaa gctctgat 89

<210> 20

<211> 89

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 20

taatctgtcc cagatccacc gccgtgaac cttgatggga ccccgagtg taaatttgat 50
gagtataaaga tcaggagctt aggggcttt 89

<210> 21

<211> 92

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 21

ggtggatctg ggacagatta tactctcacc atcagcagcc tgcagcctga agattttgca 60
acttattact gtcacagta tagtaagctt cc 92

<210> 22

<211> 80

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 22

gttttcccag tcacgaccgt acgtttaatc tctaccttgg tcccctggcc gaabgtccac 60
ggaagcttac tatactgatg 80

<210> 23

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 23

agcttccatg gacgttcggt ggaggcacca agctggaaat caaac 45

<210> 24

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 24

gtacgtttga tttccagctt ggtgcctcca ccgaacgtcc atgga 45

<210> 25

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 25

gatgcaggca gggaggatgc agtctgggt

29

<210> 26

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 26

accagactg catctctcct gcctgcac

29

<210> 27

<211> 384

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<220>

<221> CDS

<222> (1)..(384)

<400> 27

atg atg tcc tct get cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48

Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln

-20

-15

-10

-5

14 / 42

ggt acc aga tgt gac atc cag atg acc cag act gca tcc tcc ctg cct 96

Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Thr Ala Ser Ser Leu Pro

-1 1

5

10

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144

Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp

15

20

25

att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc cct 192

Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro

30

35

40

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240

Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser

45

50

55

60

agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288

Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser

65

70

75

agc ctg cag cct gaa gat ttt gca act tat tac tgt cat cag tat agt 336

Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys His Gln Tyr Ser

80

85

90

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384

Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg

95

100

105

<210> 28

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 28

actgatgaca gaaataagtt gcaaaa

26

<210> 29

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 29

ttttgcaact tattctgtc atcagt

26

<210> 30

<211> 384

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<220>

<221> CDS

<222> (1)..(384)

<400> 30

atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
 Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Leu Cys Phe Gln
 -20 -15 -10 -5

ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96
 Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
 -1 1 5 10

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
 Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
 15 20 25

att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc cct 192
 Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro
 30 35 40

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240
 Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
 45 50 55 60

agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288
 Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser
 65 70 75

agc ctg cag cct gaa gat ttt gca act tat ttc tgt cat cag tat agt 336
 Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Phe Cys His Gln Tyr Ser
 80 85 90

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384
 Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg

95

100

105

<210> 31

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 31

ggagcttaac ggctttgtct ggtttctg

28

<210> 32

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 32

cagaaaccag acaaagccgt taagctcc

28

<210> 33

<211> 384

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<220>

<221> CDS

<222> (1)..(384)

<400> 33

atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Leu Cys Phe Gln
-20 -15 -10 -5

ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96
Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
-1 1 5 10

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
15 20 25

att agt aat tat tta aac tgg tat cag cag aaa cca gac aaa gcc gtt 192
Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Asp Lys Ala Val
30 35 40

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240
Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
45 50 55 60

agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288
Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser
65 70 75

agc ctg cag cct gaa gat ttt gca act tat tac tgt cat cag tat agt 336

19 / 42

Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys His Gln Tyr Ser

80

85

90

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384

Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg

95

100

105

<210> 34

<211> 48

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 34

gttgcatat cttcaggctg cagattgctg atggtgagac tataatct 48

<210> 35

<211> 48

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 35

agattatagt ctcaccatca gcaatctgca gcctgaagat atcgcaac 48

<210> 36

<211> 384

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<220>

<221> CDS

<222> (1)..(384)

<400> 36

atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
 Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Leu Cys Phe Gln
 -20 -15 -10 -5

ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96
 Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
 -1 1 5 10

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
 Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
 15 20 25

att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc cct 192
 Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro
 30 35 40

aag ctc ctg atc ttt tac tca tca aat tta cac tgg ggg gtc cca tca 240
 Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
 45 50 55 60

agg ttc agc ggc ggt gga tct ggg aca gat tat agt ctc acc atc ago 288

Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser

65

70

75

aat ctg cag cct gaa gat atc gca act tat tac tgt cat cag tat agt 336

Asn Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys His Gln Tyr Ser

80

85

90

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384

Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg

95

100

105

<210> 37

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 37

ttcaggctgc agattgctga tgggtg

25

<210> 38

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 38

caccatcagc aatctgcagc ctgaa

25

<210> 39

<211> 384

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<220>

<221> CDS

<222> (1)..(384)

<400> 39

atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48

Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Leu Cys Phe Gln

-20

-15

-10

-5

ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96

Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser

-1 1

5

10

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144

Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp

15

20

25

att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc cct 192

Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro

30

35

40

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240

23 / 42

Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
45 50 55 60

agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288
Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser
65 70 75

aat ctg cag cct gaa gat ttt gca act tat tac tgt cat cag tat agt 336
Asn Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys His Gln Tyr Ser
80 85 90

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384
Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
95 100 105

<210> 40

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 40

gacagaaata agttgcgata tcttcaggct

30

<210> 41

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 41

agcctgaaga tategcaact tatttctgtc

30

<210> 42

<211> 384

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<220>

<221> CDS

<222> (1)..(384)

<400> 42

atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48

Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Leu Cys Phe Gln

-20 -15 -10 -5

ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96

Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser

-1 1 5 10

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144

Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp

15 20 25

att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc cct 192

25 / 42

Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro

30

35

40

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240

Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser

45

50

55

60

agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288

Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser

65

70

75

agc ctg cag cct gaa gat atc gca act tat ttc tgt cat cag tat agt 336

Ser Leu Gln Pro Glu Asp Ile Ala Thr Tyr Phe Cys His Gln Tyr Ser

80

85

90

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384

Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg

95

100

105

<210> 43

<211> 87

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 43

gcactacaag tgatggtgac tctgtctcct acagatgcag gcaggaggga tgcagactgg 60

gtcatctgga tgccacatct ggtacct

87

<210> 44

<211> 89

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 44

gtcaccatca cttgtagtgc aagtcaggac attagtaatt atttaaactg gtatcagcag 60

aaaccaggga aagccgttaa gctcctgat

89

<210> 45

<211> 89

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 45

taatctgtcc cagatccacc gccgctgaac cttgatggga cccccgagtg taaatttgat 60

gagtaaaaga tcaggagctt aacggcttt

89

<210> 46

<211> 92

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 46

ggtggatctg ggacagatta tactctcacc atcagcagcc tgcagcctga agattttgca 60
 acttatttct gtcacagta tagtaagctt cc 92

<210> 47

<211> 384

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<220>

<221> CDS

<222> (1)..(384)

<400> 47

atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
 Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
 -20 -15 -10 -5

ggt acc aga tgt gac atc cag atg acc cag tct gca tcc tcc ctg cct 96
 Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Ala Ser Ser Leu Pro
 -1 1 5 10

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
 Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
 15 20 25

att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc gtt 192
 Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Val

28 / 42

30

35

40

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240

Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser

45

50

55

60

agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288

Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser

65

70

75

agc ctg cag cct gaa gat ttt gca act tat ttc tgt cat cag tat agt 336

Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Phe Cys His Gln Tyr Ser

80

85

90

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384

Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg

95

100

105

<210> 48

<211> 384

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<220>

<221> CDS

<222> (1)..(384)

<400> 48

atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48

Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Leu Cys Phe Gln

-20

-15

-10

-5

ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96

Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser

-1 1

5

10

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144

Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp

15

20

25

att agt aat tat tta aac tgg tat cag cag aaa cca gac aaa gcc gtt 192

Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Asp Lys Ala Val

30

35

40

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240

Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser

45

50

55

60

agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288

Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser

65

70

75

agc ctg cag cct gaa gat atc gca act tat ttc tgt cat cag tat agt 336

Ser Leu Gln Pro Glu Asp Ile Ala Thr Tyr Phe Cys His Gln Tyr Ser

80

85

90

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384

Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg

<210> 49

<211> 76

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 49

catgcatgag gctctgcaca accactacac gcagaagagc ctctccctgt ctcccggggg 60
agaattcatt gatcag 76

<210> 50

<211> 85

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 50

gatactgata aatgaattct cccccgggag acagggagag gctcttctgc gtgtagtggt 60
tgtgcagagc ctcatgcatg gggcc 85

<210> 51

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 51

gtctcccgga aaagcaccta ctagtagttc tacaaag

37

<210> 52

<211> 38

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 52

ccctgatcaa tgaattcaag tcagtgttga gatgatgc

38

<210> 53

<211> 582

<212> PRT

<213> Artificial Sequence

<400> 53

Glu Val Gln Leu Val Glu Ser Gly Gly Asp Phe Val Gln Pro Gly Gly

1

5

10

15

Ser Leu Arg Val Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser His Tyr

20

25

30

Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

35

40

45

Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val

50

55

60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr

65

70

75

80

Leu Gln Met Arg Ser Leu Arg Ala Glu Asp Ser Ala Val Tyr Phe Cys

85

90

95

Thr Arg Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser Trp Gly Gln Gly

100

105

110

Thr Leu Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe

115

120

125

Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu

130

135

140

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp

145

150

155

160

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu

165

170

175

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser

180

185

190

Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro

195

200

205

Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys

210

215

220

Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro

225

230

235

240

Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser

245

250

255

Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp

260

265

270

Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn

275

280

285

Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val

290

295

300

Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu

305

310

315

320

Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys

325

330

335

Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr

340

345

350

Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr

355

360

365

Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu

370

375

380

Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu

385

390

395

400

Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys

405

410

415

Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu

420

425

430

Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly

435

440

445

Lys Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu

450

455

460

His Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr

465

470

475

480

Lys Asn Pro Lys Leu Thr Arg Met Leu Thr Phe Lys Phe Tyr Met Pro

485

490

495

Lys Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu

500

505

510

Lys Pro Leu Glu Glu Val Leu Asn Leu Ala Gln Ser Lys Asn Phe His

515

520

525

Leu Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu
 530 535 540

Leu Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr
 545 550 555 560

Ala Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser
 565 570 575

Ile Ile Ser Thr Leu Thr
 580

<210> 54

<211> 108

<212> PRT

<213> Artificial Sequence

<400> 54

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr
 20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Asp Lys Ala Val Lys Leu Leu Ile
 35 40 45

Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60

Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80

Glu Asp Ile Ala Thr Tyr Phe Cys His Gln Tyr Ser Lys Leu Pro Trp
 85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
 100 105

<210> 55

<211> 119

<212> PET

<213> Mus musculus

<400> 55

Glu Val Thr Leu Val Glu Ser Gly Gly Asp Phe Val Lys Pro Gly Gly
 1 5 10 15

Ser Leu Lys Val Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser His Tyr
 20 25 30

Ala Met Ser Trp Val Arg Gln Thr Pro Ala Lys Arg Leu Glu Trp Val
 35 40 45

Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
 65 70 75 80

37 / 42

Leu Gln Met Arg Ser Leu Arg Ser Glu Asp Ser Ala Met Tyr Phe Cys

85

90

95

Thr Arg Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser Trp Gly Gln Gly

100

105

110

Thr Thr Leu Thr Val Ser Ser

115

<210> 56

<211> 108

<212> PRT

<213> Mus musculus

<400> 56

Asp Ile Gln Met Thr Gln Thr Ala Ser Ser Leu Pro Ala Ser Leu Gly

1

5

10

15

Asp Arg Val Thr Ile Ser Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr

20

25

30

Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val Lys Leu Leu Ile

35

40

45

Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly

50

55

60

Gly Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser Asn Leu Glu Pro

65

70

75

80

Glu Asp Ile Ala Thr Tyr Phe Cys His Gln Tyr Ser Lys Leu Pro Trp

85

90

95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg

100

105

<210> 57

<211> 582

<212> PRT

<213> Artificial Sequence

<400> 57

Glu Val Thr Leu Val Glu Ser Gly Gly Asp Phe Val Lys Pro Gly Gly

1

5

10

15

Ser Leu Lys Val Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser His Tyr

20

25

30

Ala Met Ser Trp Val Arg Gln Thr Pro Ala Lys Arg Leu Glu Trp Val

35

40

45

Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val

50

55

60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr

65

70

75

80

Leu Gln Met Arg Ser Leu Arg Ser Glu Asp Ser Ala Met Tyr Phe Cys

85

90

95

Thr Arg Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser Trp Gly Gln Gly
 100 105 110

Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
 115 120 125

Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
 130 135 140

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
 145 150 155 160

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
 165 170 175

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
 180 185 190

Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro
 195 200 205

Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys
 210 215 220

Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro
 225 230 235 240

Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser
 245 250 255

40 / 42

Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp
260 265 270

Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn
275 280 285

Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val
290 295 300

Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu
305 310 315 320

Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys
325 330 335

Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr
340 345 350

Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr
355 360 365

Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu
370 375 380

Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu
385 390 395 400

Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys
405 410 415

41 / 42

Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
420 425 430

Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
435 440 445

Lys Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu
450 455 460

His Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr
465 470 475 480

Lys Asn Pro Lys Leu Thr Arg Met Leu Thr Phe Lys Phe Tyr Met Pro
485 490 495

Lys Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu
500 505 510

Lys Pro Leu Glu Glu Val Leu Asn Leu Ala Gln Ser Lys Asn Phe His
515 520 525

Leu Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu
530 535 540

Leu Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr
545 550 555 560

Ala Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser
565 570 575

... ..

42 / 42

Ile Ile Ser Thr Leu Thr

580

... ..

I, as a below named inventor, hereby declare that my residence, post office address and citizenship are as stated below next to my name, and I believe subject matter which is claimed and for which a patent is sought on the invention entitled, **GRAFTED ANTIBODY AGAINST GANGLIOSIDE GD3 AND DERIVATIVE OF ANTIBODY AGAINST GANGLIOSIDE GD3** is the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the invention.

The specification of which (check applicable box(es))

☒ was filed as PCT international application No. PCT/JP00/06774 on September 29, 2000 as U.S. Application Serial No. _____ and if applicable to U.S. or PCT application) was amended on _____

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with 37 C.F.R. 1.56. I hereby claim foreign priority benefits under 35 U.S.C. 119/365 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application(s) for patent or inventor's certificate having a filing date before that of the application on which priority is claimed or, if no priority is claimed, before the filing date of this application.

Application Number	Country	Day/Month/Year Filed
P. 2000-105088	Japan	30/September/1999
P. Hei 11-278291	Japan	6/April/2000

I hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below

I hereby claim the benefit under 35 U.S.C. 120/365 of all prior United States and PCT international applications listed above or below and, insofar as the subject matter of each of the claims of this application is not disclosed in such prior applications in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose material information as defined in 37 C.F.R. 1.56 which occurred between the filing date of the prior applications and the national or PCT international filing date of this application.

Prior U.S./PCT Application(s): _____
Application Serial No. _____
Day/Month/Year Filed _____
Status: patented _____
pending, abandoned _____

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon. And I hereby appoint **NIXON & VANDERHYE P.C., 1100 North Glebe Rd., 8th Floor, Arlington, VA 22201-4714, telephone number (703) 816-4000** (to whom all communications are to be directed), and the following attorneys thereof (of the same address) individually and collectively my attorneys to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith and with the resulting patent. Arthur R. Crawford, 25322, Larry S. Nixon, 25640, Robert A. Vanderhye, 27076, James T. Hosmer, 30184, Robert W. Farns, 31352, Richard G. Beshar, 22770, Mark E. Nusbaum, 32348, Michael J. Keenan, 32106, Bryan H. Davidson, 30251, Stanley C. Spooner, 27393, Leonard C. Mitchell, 29009, Duane M. Byers, 33363, Jeffrey H. Nelson, 30481, John R. Lastova, 33149, H. Warren Burman, Jr. 29366, Thomas E. Byrne, 32205, Mary J. Wilson, 32955, J. Scott Davidson, 33489, Alan M. Kagen, 36178, William J. Griffin, 31280, Robert A. Molan, 29834, B. J. Sadoff, 36663, James D. Benquist, 34776, Updeep S. Gill, 37334.

Inventor's Signature	Date	Inventor	Residence (city)	Post Office Address	(Zip Code)
<i>Nobuo</i>	March 25, 2002	Nobuo	Machida-shi, Tokyo	c/o Tokyo Research Laboratories, KYOWA HAKKO KOGYO CO. LTD., 6-6, Asahi-machi 3-chome,	Machida-shi, Tokyo 194-8533 Japan
<i>MI</i>		MI	(last)	(state/country)	Japan
<i>MI</i>		MI	(last)	(citizenship)	Japan
<i>Kenya</i>	March 25, 2002	Kenya	Machida-shi, Tokyo	c/o Tokyo Research Laboratories, KYOWA HAKKO KOGYO CO. LTD., 6-6, Asahi-machi 3-chome,	Machida-shi, Tokyo 194-8533 Japan
<i>MI</i>		MI	(last)	(state/country)	Japan
<i>MI</i>		MI	(last)	(citizenship)	Japan
<i>Kazuyasu</i>	March 25, 2002	Kazuyasu	Machida-shi, Tokyo	c/o Tokyo Research Laboratories, KYOWA HAKKO KOGYO CO. LTD., 6-6, Asahi-machi 3-chome,	Machida-shi, Tokyo 194-8533 Japan
<i>MI</i>		MI	(last)	(state/country)	Japan
<i>MI</i>		MI	(last)	(citizenship)	Japan

FOR ADDITIONAL INVENTORS, check box ☒ and attach sheet with same information and signature and date for each

DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION
IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Inventor's Signature: Ringer Niwa Date: March 25, 2002
Inventor: Ringer Niwa
Residence (city): Machida-shi, Tokyo
Post Office Address: c/o Tokyo Research Laboratories KYOWA HAKKO KOGYO CO., LTD., 6-6, Asahi-machi 3-chome
(Zip Code) Machida-shi, Tokyo 194-8533 Japan
Inventor's Signature: _____
Inventor: _____
Residence (city): _____
Post Office Address: _____
(Zip Code) _____

5 Inventor's Signature: _____
Inventor: _____
Residence (city): _____
Post Office Address: _____
(Zip Code) _____

6 Inventor's Signature: _____
Inventor: _____
Residence (city): _____
Post Office Address: _____
(Zip Code) _____

7 Inventor's Signature: _____
Inventor: _____
Residence (city): _____
Post Office Address: _____
(Zip Code) _____

8 Inventor's Signature: _____
Inventor: _____
Residence (city): _____
Post Office Address: _____
(Zip Code) _____

9 Inventor's Signature: _____
Inventor: _____
Residence (city): _____
Post Office Address: _____
(Zip Code) _____

10 Inventor's Signature: _____
Inventor: _____
Residence (city): _____
Post Office Address: _____
(Zip Code) _____

11 Inventor's Signature: _____
Inventor: _____
Residence (city): _____
Post Office Address: _____
(Zip Code) _____

12 Inventor's Signature: _____
Inventor: _____
Residence (city): _____
Post Office Address: _____
(Zip Code) _____